

Born-Oppenheimer Approximation

1. In the Born-Oppenheimer approximation the spatial wavefunction is approximated in the following manner:

$$\Psi(\vec{r}_e, \vec{R}_N, \sigma_e, \sigma_N) \approx \psi_{\text{electronic}}(\vec{r}_e; \vec{R}_N) \chi_{\text{nuclear}}(\vec{R}_N) \Gamma_{e\text{-spin}}(\sigma_e) \Gamma_{N\text{-spin}}(\sigma_N)$$

here $\Gamma_{e\text{-spin}}(\sigma_e) \Gamma_{N\text{-spin}}(\sigma_N)$ are the electronic and nuclear spin functions.

2. In the wavefunction $\psi_{\text{electronic}}(\vec{r}_e; \vec{R}_N)$ the electrons coordinates, \vec{r}_e , are treated as bona fide variables while the nuclear coordinates \vec{R}_N , are treated as parameters as elaborated below.
3. To get for $\psi_{\text{electronic}}(\vec{r}_e; \vec{R}_N)$, we solve the Schrodinger equation for electronic motion assuming some **fixed (relative) values** of the nuclear coordinates. For the H_2 molecule the electronic Hamiltonian is:

$$\mathcal{H}_{\text{electronic}}(\mathbf{R}_{AB}) = -\frac{1}{2} \nabla_1^2 - \frac{1}{2} \nabla_2^2 + \frac{Z_A Z_B}{R_{AB}} - \frac{1}{r_{A1}} - \frac{1}{r_{A2}} - \frac{1}{r_{B1}} - \frac{1}{r_{B2}} + \frac{1}{r_{12}}$$

where $\mathbf{R}_{AB} = |\vec{R}_A - \vec{R}_B|$, $r_{A1} = |\vec{r}_1 - \vec{R}_A| \cdots r_{12} = |\vec{r}_2 - \vec{r}_1|$

In applying this Hamiltonian a fixed value of $\mathbf{R}_{AB} \equiv R$ is selected to get

$$\mathcal{H}_{\text{el}}(R) \psi_{\text{el}}(\vec{r}_e; \vec{R}) = E(R) \psi_{\text{el}}(\vec{r}_e; \vec{R})$$

Thus we get ψ 's and E 's for various values of R . In fact when we solve the Schrodinger equation we may get several (a manifold) of wavefunctions and energies corresponding to the ground and excited states: $\psi_i(\vec{r}_e; \vec{R})$ and $E_i(R)$.

The values of $E_i(R)$ vs R give the intramolecular electronic energy for the electronic state i (the ground state or various excited states).

4. The nuclear wavefunction $\chi_{\text{nuclear}}(R)$ is determined by a Schrodinger equation for nuclear motion:

$$\mathcal{H}_{\text{nuclear}}^i \chi_{v,J}^i(R) = E_{\text{total}}^i \chi_{v,J}^i(R)$$

with $\mathcal{H}_{\text{nuclear}}^i = -\frac{1}{2\mu} \nabla_R^2 + E_{\text{el}}^i(R)$ containing the kinetic energy of relative

motion of the nuclei (rotation and vibration) and $E_{\text{el}}^i(R)$, the electronic energy of state i as a function of R , providing the potential energy for the vibrational motion of the nuclei. Note that for each electronic state i there will be a set of rotational - vibrational functions with vibrational quantum numbers v and rotational quantum numbers J .

5. The justification for the Born-Oppenheimer approximation is that electronic motion is much more rapid than nuclear motion; the electron distribution effectively 'sees' a series of fixed nuclear positions.