

# Chemistry 163B

$\mu_i$  and  $\Delta\mu_{\text{reaction}}$

Activity

Equilibrium

1

## *goals*

---


- Derive equilibrium and spontaneity criteria applying multicomponent thermodynamic relationships; i.e chemical potential ( $\Delta\mu_{\text{reaction}}$ )
- Define concentration dependence of  $\mu$  in terms of **activity** (fugacity) of 'real' gases, actual solutes
- Apply **activity** to equilibrium  $K_{\text{eq}}$
- Derive how to obtain fugacity if **REAL** gas

2

Chemistry 163B, Winter 2013  
Lecture 17- Chemical Potential and Activity

*equilibrium in terms of  $\Delta\mu$*

---



$$v_A = -n_A \quad v_B = -n_B \quad v_C = +n_C \quad v_D = +n_D$$

$d\xi$  is extent of reaction

$d\xi > 0$  forward reaction

$d\xi < 0$  reverse reaction

$$dn_i = v_i d\xi$$

$dn_i > 0$  substance  $i$  increases  
 $dn_i < 0$  substance  $i$  decreases

3

*equilibrium in terms of  $\Delta\mu$*

---

$dG_{T,P} < 0$  spontaneous

$dG_{T,P} = 0$  equilibrium

whole pot of mixed reactants and products

$$dG = -SdT + VdP + \sum_{i=1}^N \mu_i dn_i \quad dn_i = v_i d\xi$$

$$dG = -SdT + VdP + \sum_{i=1}^N \mu_i v_i d\xi$$

$$dG_{T,P} = \left( \sum_{i=1}^N \mu_i v_i \right) d\xi \leq 0$$

4

Chemistry 163B, Winter 2013  
Lecture 17- Chemical Potential and Activity

*equilibrium in terms of  $\Delta\mu$*

---

$$dG_{T,P} = \underbrace{\left( \sum_{i=1}^N \mu_i \nu_i \right)}_{?} d\xi \leq 0$$

$$dG_{T,P} = \underbrace{\left( \sum_{i=1}^N \mu_i \nu_i \right)}_{\Delta\mu_{\text{reaction}}} d\xi \leq 0$$

$$dG_{T,P} = \Delta\mu_{\text{reaction}} d\xi \leq 0$$

$\Delta\mu_{\text{reaction}} < 0$  forward reaction spontaneous ( $d\xi > 0$ )

$\Delta\mu_{\text{reaction}} > 0$  reverse reaction spontaneous ( $d\xi < 0$ )

$\Delta\mu_{\text{reaction}} = 0$  equilibrium

just like  $\Delta G$  !!!

5

*concentration dependence of  $\mu_i$*

---

ideal gas, one component (pure substance)

$$\bar{G} = \bar{G}^\circ + RT \ln \left( \frac{P}{1 \text{ bar}} \right)$$

led to

$$\Delta G_{\text{reaction}} = \Delta G_{\text{reaction}}^\circ + RT \ln(Q_P)$$

what about if other species present?

$$\mu_i = \mu_i^\circ + RT \ln \left( \frac{P_i}{1 \text{ bar}} \right)$$

$$\Delta\mu_{\text{reaction}} = \Delta\mu_{\text{reaction}}^\circ + RT \ln Q_P$$

**HANDOUT #45**

$$\Delta\mu_{\text{reaction}}^\circ = \sum_i \nu_i \mu_i^\circ \quad Q_P = \prod_i \left( \frac{P_i}{1 \text{ bar}} \right)^{\nu_i}$$

6

Chemistry 163B, Winter 2013  
Lecture 17- Chemical Potential and Activity

*yada -yada- yada: and so forth for  $\Delta\mu_{\text{reaction}}$*


$$\Delta\mu_{\text{reaction}}^{\circ} = -RT \ln K_P$$

$$\left( \frac{\partial \frac{\Delta\mu}{T}}{\partial T} \right)_P = -\frac{\Delta H}{T^2} \quad \text{where} \quad \Delta H = \sum_i \nu_i \bar{H}_i = \sum_i \nu_i \left( \frac{\partial H}{\partial n_i} \right)_{T,P,n_j \neq n_i}$$

$$\left( \frac{\partial \ln K}{\partial T} \right)_P = \frac{\Delta H^{\circ}}{RT^2} \quad \text{where} \quad \Delta H^{\circ} = \sum_i \nu_i \bar{H}_i^{\circ}$$

7

*correcting for REALITY (activity and fugacity)*



I STILL REFUSE  
TO ACCEPT REALITY



My childhood dream  
was to become a mermaid  
with a blue tail

8

Chemistry 163B, Winter 2013  
Lecture 17- Chemical Potential and Activity

*correcting for REALITY (activity and fugacity)*

---

- will define activity and fugacity coefficients  $\gamma_i$ 's that provides corrections for the deviation of chemical potential from the **ideal** gas and solute concentration dependence
- activity and fugacity coefficients are obtained from experimental measurements on **REAL** systems or by theory (Debye-Huckel)

9

*more general  $\mu_i$  and corrections for non-ideality (~E&R eqn 9.50, p227)<sub>3rd</sub>*

---

$$\mu_i(T) = \mu_i^\circ(T) + RT \ln \left( \frac{a_i}{a_i^\circ} \right)$$

standard conditions

how  $\mu_i$  'really' changes in going from standard conditions to actual conditions

$a_i$  = activity of component  $i$

$a_i = \gamma_i \times [\text{ideal measure of pressure, concentration, etc}]$

$\gamma_i$  is activity coefficient, a correction for non-ideality

$a_i^\circ = 1$  unit (bar, molar, etc)

10

Chemistry 163B, Winter 2013  
Lecture 17- Chemical Potential and Activity

more general  $\mu_i$  and corrections for non-ideality

$$\Delta\mu_{\text{reaction}} = \Delta\mu^\circ + RT \ln Q$$

where now  $Q$  is written in terms of activities

$$Q = \prod_i \underbrace{\left( \frac{a_i}{a_i^\circ} \right)^{\bar{\nu}_i}}_{\text{unitless}} = \frac{(a_C)^{\bar{n}_C} (a_D)^{\bar{n}_D}}{(a_A)^{\bar{n}_A} (a_B)^{\bar{n}_B}}$$

dropped the  $a^\circ=1$  'unit'  
but  $Q$  is 'unitless'

11

fugacity of gases

1. GASES:

$$a_i = f_i = \gamma_i P_i$$

partial pressure

fugacity of gas

fugacity coefficient

$$Q = \frac{(a_C)^{\bar{n}_C} (a_D)^{\bar{n}_D}}{(a_A)^{\bar{n}_A} (a_B)^{\bar{n}_B}}$$

$$Q = \frac{\left( \gamma_C \frac{P_C}{1\text{bar}} \right)^{\bar{n}_C} \left( \gamma_D \frac{P_D}{1\text{bar}} \right)^{\bar{n}_D}}{\left( \gamma_A \frac{P_A}{1\text{bar}} \right)^{\bar{n}_A} \left( \gamma_B \frac{P_B}{1\text{bar}} \right)^{\bar{n}_B}}$$

correction for non-ideality

$$Q = \frac{\left( \frac{\gamma_C^{\bar{n}_C} \gamma_D^{\bar{n}_D}}{\gamma_A^{\bar{n}_A} \gamma_B^{\bar{n}_B}} \right) \left( \frac{P_C}{1\text{bar}} \right)^{\bar{n}_C} \left( \frac{P_D}{1\text{bar}} \right)^{\bar{n}_D}}{\left( \frac{P_A}{1\text{bar}} \right)^{\bar{n}_A} \left( \frac{P_B}{1\text{bar}} \right)^{\bar{n}_B}} = \gamma Q_P$$

ideal gas  $Q_P$

12

Chemistry 163B, Winter 2013  
Lecture 17- Chemical Potential and Activity

*other conventions for activities*

---

2. pure solids and liquids

$$\mu_i(T, P) \approx \mu_i^\circ(T, P = 1 \text{ bar})$$

$$\left( \frac{\partial \mu_i}{\partial P} \right)_T = \bar{V}_i \quad (\text{small for liquid or solid})$$

so  $a_i \approx 1$  for pure solid or liquid  
[unless extreme pressure]

13

*other conventions for activities*

---

3. solutes in solutions

$$a_i = \gamma_i [I]$$

concentration of I,  
usually molar  
but may be  $X_i$

activity coefficient  $\gamma_i$  corrects 'ideal' measure of 'concentration'

if "activity coefficients unity"

$$a_i = [I] \quad a_i \equiv f_i = P_i \quad a_i = 1$$


*solute*      *gas*      *pure liquid or solid*

HW#7     $\gamma=1$  except prob. 41\* and 43.

14

Chemistry 163B, Winter 2013  
Lecture 17- Chemical Potential and Activity

how to evaluate activity (fugacity) coefficients for real gases  
(HW7 #\*41)



15

how to evaluate activity (fugacity) coefficients for real gases

$$\left(\frac{\partial \mu}{\partial P}\right)_{T,n} = \bar{V} \quad \text{and}$$

$$\mu = \mu^\circ + RT \ln f$$

need in a moment

$$\left(\frac{\partial \mu}{\partial P}\right)_{T,n} = RT \left(\frac{\partial \ln f}{\partial P}\right)_{T,n}$$

$$\text{so } RT \left(\frac{\partial \ln f}{\partial P}\right)_{T,n} = \bar{V}$$

expression for  $d \ln \left(\frac{f}{P}\right)$  will prove useful

$$\begin{aligned} \left(\frac{\partial \ln \left(\frac{f}{P}\right)}{\partial P}\right)_{T,n} &= \left(\frac{\partial (\ln f - \ln P)}{\partial P}\right)_{T,n} = \left(\frac{\partial \ln f}{\partial P}\right)_{T,n} - \left(\frac{\partial \ln P}{\partial P}\right)_{T,n} \\ &= \frac{1}{RT} \bar{V} - \frac{1}{P} = \frac{1}{RT} \left(\bar{V} - \frac{RT}{P}\right) \end{aligned}$$

16



Chemistry 163B, Winter 2013  
Lecture 17- Chemical Potential and Activity

how to evaluate activity (fugacity) coefficients for real gases

$$\int_{P_1}^P d \left( \ln \frac{f}{P'} \right) = \int_{P_1}^P \frac{1}{RT} \left( \bar{V} - \frac{RT}{P'} \right) dP'$$

$$\ln \left( \frac{f(P)}{P} \right) - \ln \left( \frac{f(P_1)}{P_1} \right) = \frac{1}{RT} \int_{P_1}^P \left( \bar{V} - \frac{RT}{P'} \right) dP'$$

$$\ln \left( \frac{f(P)}{P} \right) = \ln \left( \frac{f(P_1)}{P_1} \right) + \frac{1}{RT} \int_{P_1}^P \left( \bar{V} - \frac{RT}{P'} \right) dP'$$

$$P_1 \rightarrow 0$$

$$\frac{f(P_1)}{P_1} \rightarrow 1$$

$$\ln \left( \frac{f(P_1)}{P_1} \right) \rightarrow 0$$

$$(\bar{V}_{ACTUAL} - \bar{V}_{IDEAL GAS})$$

17

how to evaluate activity (fugacity) coefficients for real gases  
(eqn 7.20 E&R and HW7 #41)

$$\ln f(P) = \ln P + \frac{1}{RT} \int_{P_1 \rightarrow 0}^P \left( \bar{V} - \frac{RT}{P'} \right) dP' = \ln P + \frac{1}{RT} \int_{P_1 \rightarrow 0}^P (\bar{V}_{ACTUAL} - \bar{V}_{IDEAL GAS}) dP'$$

$$\ln \left( \frac{f(P)}{P} \right) = \ln(\gamma) = \frac{1}{RT} \int_{P_1 \rightarrow 0}^P \left( \bar{V} - \frac{RT}{P'} \right) dP' = \frac{1}{RT} \int_{P_1 \rightarrow 0}^P (\bar{V}_{ACTUAL} - \bar{V}_{IDEAL}) dP'$$

$$z = \frac{\bar{V}_{actual}}{\bar{V}_{ideal}} = \frac{P\bar{V}_{actual}}{RT} \quad (\text{compression factor E\&R eqn. 7.6})$$

$$\ln \gamma = \frac{1}{RT} \int_{P_1 \rightarrow 0}^P \bar{V}_{ideal} (z-1) dP' = \int_{P_1 \rightarrow 0}^P \frac{(z-1)}{P'} dP' \quad \text{HW7 41*}$$

$$\gamma(P, T) = \exp \left[ \int_{P_1 \rightarrow 0}^P \frac{z-1}{P'} dP' \right] \quad (\text{E \& R eqn 7.21})$$

18

*End of Lecture*