

Chemistry 163B
Colligative Properties
Challenged Penpersonship
Notes

1

colligative properties of solutions



colligative 

One entry found.

Main Entry: col·li·ga·tive

Pronunciation: 'kă-lə-gā-tiv, kə-'li-gə-tiv

Function: *adjective*

: depending on the number of particles (as molecules) and not on the nature of the particles
<pressure is a *colligative property*>

<http://www.merriam-webster.com/dictionary/colligative>

2

quantitative treatment of colligative properties

Handout #52

- A. Freezing point depression**
- B. Boiling Point Elevation**
- C. Osmotic Pressure**

3

quantitative treatment of colligative properties

- I. The pure solvent (component B) is originally in equilibrium in the two phases.
- II. Addition of solute (component A) lowers the chemical potential of the solvent in the solution phase
- III. Temperature (freezing point depression, boiling point elevation) or pressure (osmotic pressure) must be altered to reestablish equilibrium between the solution and the pure solvent phase.
- IV. Obtain relationships between X_A or X_B and change in T or P.

4

freezing point depression (solid \rightleftharpoons solution)

I. pure solvent is originally in equilibrium in the two phases

*pure solid*_B^{*} \rightleftharpoons *pure liquid*_B^{*} at T_f^* the normal melting T_{fusion}

$$\mu_B^{s*}(T_f^*) = \mu_B^{\ell*}(T_f^*)$$

$$\Delta\mu_B(T_f^*) = \mu_B^{\ell*}(T_f^*) - \mu_B^{s*}(T_f^*) = 0$$

$$\Delta\bar{H}(T_f^*) = \Delta\bar{H}_{B \text{ melting}} > 0 \quad \text{for solid} \rightarrow \text{liquid}$$

5

freezing point depression (solid \rightleftharpoons solution)

II. Still at T_f^* , add solute A to solvent with resulting mole fractions X_A and X_B

for solid phase of B there is no change :

$$\mu_B^{s*}(T_f^*) = \mu_B^{solid}(T_f^*)$$

for the solvent (B) in solution:

$$\mu_B^{\ell}(T_f^*) \equiv \mu_B^{solvent} \equiv \mu_B^{\ell(in \text{ soln})}(T_f^*) = \mu_B^{\ell*}(T_f^*) + RT_f^* \ln(\gamma_B X_B)$$

$$\text{so now } \Delta\mu_B(T_f^*) = \mu_B^{\ell}(T_f^*) - \mu_B^{s*}(T_f^*) = \Delta\mu_B^*(T_f^*) + RT_f^* \ln(\gamma_B X_B)$$

$$\text{where } \Delta\mu_B^*(T_f^*) = \mu_B^{\ell*}(T_f^*) - \mu_B^{s*}(T_f^*)$$

and $\Delta\mu_B^*(T_f^*) = 0$ since pure liquid and solid are in equilibrium at T_f^*

$$\text{thus } \Delta\mu_B(T_f^*) = RT_f^* \ln(\gamma_B X_B) < 0$$

so the forward reacton (melting of the solid)
would now occur spontaneously at T_f^*

6



freezing point depression (solid \rightleftharpoons solution)

III. Alter temperature to restore equilibrium $T_f^\circ \rightarrow T_f$

$$\left(\frac{\partial \frac{\Delta\mu}{T}}{\partial T} \right)_P = -\frac{\Delta\bar{H}}{T^2}$$

$$\int_{T_f^\circ}^{T_f} d\left(\frac{\Delta\mu_B}{T} \right)_P = -\int_{T_f^\circ}^{T_f} \frac{\Delta\bar{H}_{B \text{ melting}}}{T^2} dT$$

old stuff

$$\left(\frac{\Delta\mu_B(T_f)}{T_f} \right)_P - \left(\frac{\Delta\mu_B(T_f^\circ)}{T_f^\circ} \right)_P = -\int_{T_f^\circ}^{T_f} \frac{\Delta\bar{H}_{B \text{ melting}}}{T^2} dT$$

7

freezing point depression (solid \rightleftharpoons solution)

III. Alter temperature to restore equilibrium (continued)

$$\left(\frac{\Delta\mu_B(T_f)}{T_f} \right)_P - \left(\frac{\Delta\mu_B(T_f^\circ)}{T_f^\circ} \right)_P = -\int_{T_f^\circ}^{T_f} \frac{\Delta\bar{H}_{B \text{ melting}}}{T^2} dT$$

$$\left(\frac{\Delta\mu_B(T_f)}{T_f} \right)_P = 0 \text{ since at 'new' equilibrium } T_f, \Delta\mu_B(T_f) = 0$$

$$\text{and } \left(\frac{\Delta\mu_B(T_f^\circ)}{T_f^\circ} \right)_P = R \ln(\gamma_B X_B) \text{ from eqn in II.}$$

$$-R \ln(\gamma_B X_B) = -\int_{T_f^\circ}^{T_f} \frac{\Delta\bar{H}_{B \text{ melting}}}{T^2} dT$$

$$R \ln(\gamma_B X_B) + \left[-\int_{T_f^\circ}^{T_f} \frac{\Delta\bar{H}_{B \text{ melting}}}{T^2} dT \right] = 0$$

change in $\Delta\mu_B$ due to adding solute

change in $\Delta\mu_B$ due to temperature change

8

freezing point lowering

IV. Obtain relationships between X_B and change in T

$$R \ln(\gamma_B X_B) = \int_{T_f^*}^{T_f} \frac{\Delta \bar{H}_{B \text{ melting}}}{T^2} dT$$

$\Delta \bar{H}_{B \text{ melting}} \sim$ independent of T

$$R \ln(\gamma_B X_B) = -\Delta \bar{H}_{B \text{ melting}} \left[\frac{1}{T_f} - \frac{1}{T_f^*} \right]$$

since $lfs < 0 \Rightarrow T_f < T_f^*$ (freezing point **depression**)

$$\gamma_B X_B = \exp \left[-\frac{\Delta \bar{H}_{B \text{ melting}}}{R} \left[\frac{1}{T_f} - \frac{1}{T_f^*} \right] \right] \quad (\text{integration of eqn 9.31 E\&R})$$

9

freezing point lowering

IV. Obtain relationships between X_B and change in T (*cont*)


$$\gamma_B X_B = \exp \left[-\frac{\Delta \bar{H}_{B \text{ melting}}}{R} \left[\frac{1}{T_f} - \frac{1}{T_f^*} \right] \right]$$

$$-\frac{R}{\Delta \bar{H}_{B \text{ melting}}} \ln(\gamma_B X_B) + \frac{1}{T_f^*} = \frac{1}{T_f}$$

$$T_f = \frac{T_f^* \Delta \bar{H}_{B \text{ melting}}}{\Delta \bar{H}_{B \text{ melting}} - RT_f^* \ln(\gamma_B X_B)} \quad (\sim \text{eqn 9.32 E\&R})$$

10


Osmotic Pressure Equilibrium



11

osmotic pressure (pure solvent \rightleftharpoons solution [solvent + solute])

- I. pure solvent at P_{left} is originally in equilibrium with pure solvent at P_{right} ; i.e. $P_{\text{left}} = P_{\text{right}} = P_0$



'left' and 'right' refer to compartments separated by solute impermeable membrane

$$\mu_B^*(P_0, \text{left}) = \mu_B^*(P_0, \text{right})$$

12

osmotic pressure (II add solute to left compartment)

II. in left hand compartment add solute A to solvent
with resulting mole fractions X_A and X_B

add X_A solute to liquid in 'left' compartment resulting in X_B for solvent

$$\mu_B^l(P_0, \text{left}) = \mu_B^{\text{pure}}(P_0, \text{left}) + RT \ln(\gamma_B X_B) < \mu_B^{\text{pure}}(P_0, \text{right})$$

$$\mu_B^l(P_0, \text{left}) < \mu_B^{\text{pure}}(P_0, \text{right})$$

so the solvent B moves spontaneously left ← right

(i.e. diluting solution on left)

13

osmotic pressure (III, alter pressure)

III. alter Pressure: $P_{\text{left}} \rightarrow P_0 + \pi$ to restore equilibrium

solution ($X_B, P_0 + \pi, \text{left}$) \rightleftharpoons pure solvent (P_0, right)

$$\left(\frac{\partial \mu_B^{\text{left}}}{\partial P} \right)_T = \bar{V}_B$$

$$\int_{P_0}^{P_0 + \pi} d\mu_B^{\text{left}}(X_B) = \int_{P_0}^{P_0 + \pi} \bar{V}_B dP$$

assuming solvent is incompressible

(\bar{V}_B doesn't change with pressure at constant T)

$$\mu_B^{\text{left}}(X_B, P_0 + \pi) = \mu_B^{\text{left}}(X_B, P_0) + [P_0 + \pi - P_0] \bar{V}_B$$

$$\mu_B^{\text{left}}(X_B, P_0 + \pi) = \mu_B^{\text{left}}(X_B, P_0) + \pi \bar{V}_B$$

14

Chemistry 163B, Winter 2013
Lecture 23 Colligative Properties

osmotic pressure (III, alter pressure, continued)

$$\mu_B^{\text{left}}(X_B, P_0 + \pi) = \mu_B^{\text{left}}(X_B, P_0) + \pi \bar{V}_B$$

$$\mu_B^{\text{left}}(X_B, P_0 + \pi) = \mu_B^{\bullet}(P_0) + RT \ln(\gamma_B X_B) + \pi \bar{V}_B$$

want π to restore equilibrium such that

$$\mu_B^{\text{left}}(X_B, P_0 + \pi) = \mu_B^{\text{right}}(P_0)$$

$$\underbrace{\mu_B^{\bullet}(P_0) + RT \ln(\gamma_B X_B) + \pi \bar{V}_B}_{\text{left}} = \underbrace{\mu_B^{\bullet}(P_0)}_{\text{right}}$$

$$\pi = -\frac{RT \ln(\gamma_B X_B)}{\bar{V}_B}$$

15

osmotic pressure (a little more manipulation)

$$\pi = -\frac{RT \ln(\gamma_B X_B)}{\bar{V}_B}$$

for $\gamma_B \approx 1$ and $X_B = 1 - X_A$

$$\pi = -\frac{RT \ln(1 - X_A)}{\bar{V}_B}$$

$\ln(1 + x) \approx x$ for small x (i.e. dilute solution, X_A small)

$$\pi = \frac{X_A RT}{\bar{V}_B}$$

$X_A = \frac{n_A}{n_A + n_B}$ and $n_A + n_B \approx n_B$ for dilute solution

$$\pi \approx \frac{n_A RT}{n_B \bar{V}_B}$$

$$\pi \bar{V}_B = n_A RT$$

$$\pi V_{\text{solution}} = n_{\text{solute}} RT$$

16

quantitative treatment of colligative properties

Handout #52 Colligative Properties


from relationships for Chem 163B final:

Colligative properties:

- freezing point lowering: $\gamma_B X_B = \exp \left[-\frac{\Delta \bar{H}_{\text{fusion}}}{R} \left[\frac{1}{T_f} - \frac{1}{T_f^*} \right] \right]$
- boiling point elevation: $\gamma_B X_B = \exp \left[\frac{\Delta \bar{H}_{\text{vaporization}}}{R} \left[\frac{1}{T_b} - \frac{1}{T_b^*} \right] \right]$
- osmotic pressure: $\pi = \frac{-RT \ln(\gamma_B X_B)}{\bar{V}_B}$
 $\pi \approx \frac{n_A RT}{V_B} = \frac{n_{\text{solute}} RT}{V_{\text{solvent}}}$ for dilute solution

17

reverse osmosis




Water Systems
Aqua Technology
For the 21st Century

ESP Water Products.com

18


Reverse Osmosis



http://www.zenon.com/image/resources/glossary/reverse_osmosis/normal_osmosis.jpg

19

effect of osmosis on blood cells



20

Chemistry 163B, Winter 2013
Lecture 23 Colligative Properties

hyponatremia

Woman dies after water-drinking contest

Water intoxication eyed in 'Hold Your Wee for a Wii' contest death

AP Associated Press
updated 7:24 p.m. PT, Sat., Jan. 13, 2007

SACRAMENTO, Calif. - A woman who competed in a radio station's contest to see how much water she could drink without going to the bathroom died of water intoxication, the coroner's office said Saturday.

Jennifer Strange, 28, was found dead Friday in her suburban Rancho Cordova home hours after taking part in the "Hold Your Wee for a Wii" contest in which KDND 107.9 promised a Nintendo Wii video game system for the winner.

"She said to one of our supervisors that she was on her way home and her head was hurting her real bad," said Laura Rios, one of Strange's co-workers at Radiological Associates of Sacramento. "She was crying and that was the last that anyone had heard from her."



Woman in water drinking contest dies
Jan. 15. Sacramento Bee reporter Christina Jewett talks to MSNBC-TV's Contessa Brewer about the death of a woman who had competed in a radio station contest.
MSNBC

21

End of Lecture