

# Chemistry 163B, Winter 2014

## Lecture 16 Multicomponent Systems and Partial Molar Quantities

for save as PDF

### Chemistry 163B

### Introduction to

### Multicomponent Systems

### and Partial Molar Quantities

1

#### the problem of partial mmolar quantities

---

mix: 10 moles ethanol C<sub>2</sub>H<sub>5</sub>OH (580 mL)  
 with  
 1 mole water H<sub>2</sub>O (18 mL)  
 get (580+18)=598 mL of solution?  
 no only 594 mL

for pure H<sub>2</sub>O  $\left(\frac{\partial V}{\partial n_{H_2O}}\right)_{T,P,n_{EtOH}} = V_{H_2O}^* = 18 \text{ mL}$

but in 10 mol EtOH  $\left(\frac{\partial V}{\partial n_{H_2O}}\right)_{T,P,n_{EtOH}} = 14 \text{ mL}$

2

#### partial molar quantities (systems of variable composition)

---

system of n<sub>1</sub> moles substance 1, n<sub>2</sub> moles substance 2, ...  
 Ω some extensive property of system (volume, free energy, etc)

$$\bar{\Omega}_i = \left(\frac{\partial \Omega_{total}}{\partial n_i}\right)_{T,P,n_j \neq n_i}$$

"partial molar Ω" for component i  
 contribution of substance i to property Ω at T, P  
 when other components present at concentrations n<sub>j</sub>  
 "molar Ω" in presence of other species

3

slides 4-7 are taken from:  
[http://www.chem.ut.edu/faculty/cooke/3510/3510\\_chap7.ppt](http://www.chem.ut.edu/faculty/cooke/3510/3510_chap7.ppt)  
*apparently no longer available*

A site from: Stephen A. Cooke, Ph.D.  
 Department of Chemistry  
 University of North Texas

4

### PARTIAL MOLAR QUANTITIES

In a system that contains at least two substances, the total value of any extensive property of the system is the sum of the contribution of each substance to that property.

The contribution of one mole of a substance to the volume of a mixture is called the **partial molar volume** of that component.

$$V = f(p, T, n_A, n_B, \dots)$$

At constant T and p

$$dV = \left(\frac{\partial V}{\partial n_A}\right) dn_A + \left(\frac{\partial V}{\partial n_B}\right) dn_B + \dots$$

$$V_A^* = \left(\frac{\partial V}{\partial n_A}\right)_{p,T,n_{\neq A}}$$

5

### PARTIAL MOLAR VOLUME

Add n<sub>A</sub> of A to mixture

Very Large Mixture of A and B

Composition remains essentially unchanged. In this case:

$$V_A^* = \left(\frac{\partial V}{\partial n_A}\right)_{p,T,n_{\neq A}}$$

can be considered constant and the volume change of the mixture is n<sub>A</sub>V<sub>A</sub>. Likewise for addition of B.

The total change in volume is n<sub>A</sub>V<sub>A</sub> + n<sub>B</sub>V<sub>B</sub>. (Composition is essentially unchanged).

Scoop out of the reservoir a sample containing n<sub>A</sub> of A and n<sub>B</sub> of B its volume is n<sub>A</sub>V<sub>A</sub> + n<sub>B</sub>V<sub>B</sub>. Because V is a state function:

$$V = V_A^* n_A + V_B^* n_B + \dots$$

6

# Chemistry 163B, Winter 2014

## Lecture 16 Multicomponent Systems and Partial Molar Quantities

### PARTIAL MOLAR VOLUME

Illustration:

What is the change in volume of adding 1 mol of water to a large volume of water?

$$\text{The change in volume is } 18\text{cm}^3 \quad V_{\text{H}_2\text{O}} = \left( \frac{\partial V}{\partial n_{\text{H}_2\text{O}}} \right)_{P,T} = 18\text{cm}^3$$


A different answer is obtained if we add 1 mol of water to a large volume of ethanol.

$$\text{The change in volume is } 14\text{cm}^3 \quad V_{\text{H}_2\text{O}} = \left( \frac{\partial V}{\partial n_{\text{H}_2\text{O}}} \right)_{P,T,n(\text{C}_2\text{H}_5\text{OH})} = 14\text{cm}^3$$

7

### PARTIAL MOLAR QUANTITIES


$V_A$  is not generally a constant; it is a function of composition:



8

### Gibbs-Duhem (later)

$$X_{\text{ROH}} \left( \frac{\partial V_{\text{ROH}}}{\partial n_{\text{ROH}}} \right)_{T,P,n_{\text{H}_2\text{O}}} = -X_{\text{H}_2\text{O}} \left( \frac{\partial V_{\text{H}_2\text{O}}}{\partial n_{\text{ROH}}} \right)_{T,P,n_{\text{H}_2\text{O}}}$$



[http://www.chem.unt.edu/faculty/cooker/3510/3510\\_chap7.ppt](http://www.chem.unt.edu/faculty/cooker/3510/3510_chap7.ppt)

9

### partial molar quantities in biology

Proc. Natl. Acad. Sci. USA  
Vol. 84, pp. 7938-7942, November 1987  
Biophysics

#### The partial molar volume of water in biological membranes

(phosphatidylcholine/biurea/egg phosphatidylcholine) (phosphatidylcholine/biurea/egg phosphatidylcholine)  
JAMES R. SCHERER\*  
Western Regional Research Center, Agricultural Research Service, U.S. Department of Agriculture, Albany, CA 94718  
Communicated by Bryce Crawford, Jr., July 26, 1987

**ABSTRACT** A new algorithm is presented for interpreting the hydration dependence of x-ray diffraction measurements. The method assumes that the volume of the hydrocarbon phase of the lipid bilayer is not affected by hydration and that the volume expansion between bilayers at maximum hydration is caused by incorporation of water molecules whose partial molar volume is that of pure bulk water. These simple assumptions lead to a determination of the area expansion (and hence change in hydrocarbon-phase thickness) as a function of hydration. An analysis is made of x-ray data of the  $L_2$  and  $L_3$  phases of dimyristoyl phosphatidylcholine and the  $L_2$  phase of egg phosphatidylcholine. The partial molar volume of water depends critically on the degree of lipid hydration and the presence of tails between the head groups of adjacent lipids. The calculated head-group spacings at minimum hydration are consistent with those obtained from neutron diffraction and indicate that the methyl groups of the choline are almost in contact with corresponding groups in the opposing bilayer, contact with corresponding groups in the opposing bilayer. This calls into question the origin of the repulsive forces observed in dehydration experiments.

across the lipid part of the bilayer, and  $d_w$ , the distance across the aqueous region between bilayers.

$$d = d_0 + d_w \quad (1)$$

$d_0$  is usually obtained from  $d$  and the volume fraction,  $\phi$ :  $d_0 = d\phi$ .

$$\phi = [1 + (1 - \rho_w/\rho_l)V_w]^{-1} \quad (2)$$

where  $\rho_l$  is the weight fraction of lipid,  $\rho_w$  and  $\rho_l$  are the partial specific volumes of lipid and water (8). It is commonly assumed that  $\rho_l$  and  $\rho_w$  are both  $1 \text{ cm}^3/\text{g}$ . White and King (7) have questioned the last assumption by pointing out that it leads to a value of  $d_0$  for egg phosphatidylcholine (E-PC) in the  $L_2$  phase that changes by 1.5 Å over its full hydration range. However, the distance between phosphate groups across the bilayer,  $d_{\text{ph}}$ , obtained by their analysis of x-ray data of Torbet and Wilson (8), does not change by more than 4 Å. Sankar, Janak et al. (9) have shown that in the  $L_2$  phase of dimyristoyl phosphatidylcholine (DM-PC) at

10

### partial molar factoids #1 total differentials

- state function differentials for systems of variable composition (still  $\sigma_{\text{wofth}}=0$ )

$$U(S, V, n_1, \dots, n_N) \quad dU = TdS - PdV + \sum_{i=1}^N \left( \frac{\partial U}{\partial n_i} \right)_{S, V, n_{j \neq i}} dn_i$$

$$H(S, P, n_1, \dots, n_N) \quad dH = TdS + VdP + \sum_{i=1}^N \left( \frac{\partial H}{\partial n_i} \right)_{S, P, n_{j \neq i}} dn_i$$

$$A(T, V, n_1, \dots, n_N) \quad dA = -SdT - PdV + \sum_{i=1}^N \left( \frac{\partial A}{\partial n_i} \right)_{T, V, n_{j \neq i}} dn_i$$

$$G(T, P, n_1, \dots, n_N) \quad dG = -SdT + VdP + \sum_{i=1}^N \left( \frac{\partial G}{\partial n_i} \right)_{T, P, n_{j \neq i}} dn_i$$

11

### partial molar factoids #2 the chemical potential

- The partial molar Gibbs free energy, the chemical potential, plays a central role

$$\bar{G}_i = \left( \frac{\partial G}{\partial n_i} \right)_{T, P, n_{j \neq i}} \equiv \mu_i$$

thus

$$dG = -SdT + VdP + \sum_{i=1}^N \mu_i dn_i$$

and a very cute derivation give (see handout):

$$\mu_i = \left( \frac{\partial G}{\partial n_i} \right)_{T, P, n_{j \neq i}} = \left( \frac{\partial A}{\partial n_i} \right)_{T, V, n_{j \neq i}} = \left( \frac{\partial H}{\partial n_i} \right)_{S, P, n_{j \neq i}} = \left( \frac{\partial U}{\partial n_i} \right)_{S, V, n_{j \neq i}}$$

note: for A, H, U these are NOT partial molar quantities  $\bar{A}_i, \bar{H}_i, \text{ and } \bar{U}_i$

12

# Chemistry 163B, Winter 2014

## Lecture 16 Multicomponent Systems and Partial Molar Quantities

### factoid #3: properties of a system are sum of partial molar properties

3. An extensive property of a multi-component system is the sum of partial molar contributions from each of the components

$$V_{total} = \sum_i^N n_i \bar{V}_i = n_1 \bar{V}_1 + n_2 \bar{V}_2 + \dots$$

$$G = \sum_i^N n_i \bar{G}_i$$

$$H = \sum_i^N n_i \bar{H}_i \quad \text{note: } \bar{H}_i = \left( \frac{\partial H}{\partial n_i} \right)_{T, P, n_j, n_m} = \left( \frac{\partial H}{\partial n_i} \right)_{S, P, n_j, n_m} = \mu_i$$

etc.

13

### factoid #4: relationships among partial molar quantities

4. Relationships among thermodynamic quantities derived for one-component systems often hold for partial molar quantities

examples:

$$G \equiv H - TS \Rightarrow \bar{G}_i = \bar{H}_i - T\bar{S}_i$$

or

$$H \equiv U + PV \Rightarrow \bar{H}_i = \bar{U}_i + P\bar{V}_i$$

[proof in class for G; students do similar proof for H]

14

### factoid #5: Gibbs Duhem

5. The Gibbs-Duhem relationship shows that partial molar quantities for substances in a mixture can not vary independently

example:  $\bar{V}_i$  for a two component mixture e.g. EtOH + H<sub>2</sub>O

$$X_A \left( \frac{\partial \bar{V}_A}{\partial n_B} \right)_{T, P, n_A} = -X_B \left( \frac{\partial \bar{V}_B}{\partial n_A} \right)_{T, P, n_B}$$

$$X_{H_2O} \left( \frac{\partial \bar{V}_{H_2O}}{\partial n_{EtOH}} \right)_{T, P, n_{H_2O}} = -X_{EtOH} \left( \frac{\partial \bar{V}_{EtOH}}{\partial n_{H_2O}} \right)_{T, P, n_{EtOH}}$$


[note: the variation is with respect to one of the components  
( $\partial n_{EtOH}$  in both denominators)]

[derivation done in class]

15

### Gibbs-Duhem (slope of partial molar volume vs mole fraction)

$$\left( \frac{\partial \bar{V}_{EtOH}}{\partial n_{EtOH}} \right)_{T, P, n_{H_2O}} = - \frac{X_{H_2O}}{X_{EtOH}} \left( \frac{\partial \bar{V}_{H_2O}}{\partial n_{EtOH}} \right)_{T, P, n_{H_2O}} \quad \left( \frac{\partial \bar{V}_{H_2O}}{\partial n_{EtOH}} \right)_{T, P, n_{H_2O}} = - \frac{X_{EtOH}}{X_{H_2O}} \left( \frac{\partial \bar{V}_{EtOH}}{\partial n_{EtOH}} \right)_{T, P, n_{H_2O}}$$



$X_{EtOH} = 0$

$X_{H_2O} = 0$

[http://www.chem.unt.edu/faculty/cooke/3510/3510\\_chap7.ppt](http://www.chem.unt.edu/faculty/cooke/3510/3510_chap7.ppt)

16

*End of Lecture*

17