

# Chemistry 163B Winter 2014

## notes for lecture 5

Chemistry 163B  
Lecture 5 Winter 2014  
Challenged Penmanship  
Notes

1

total differential (math handout #4; E&R ch. 3)

infinitesimal change in value of state function (well behaved function)

$f(x, y)$  a well behaved function

$$\text{total change in } f \rightarrow df = \left(\frac{\partial f}{\partial x}\right)_y dx + \left(\frac{\partial f}{\partial y}\right)_x dy$$

2

differential of product (product rule)

$$d(xy) = ydx + xdy$$

3

example of implication of total differentials

First Law

$$dU_{sys} = dq_{sys} + \delta w_{sys} + dn_{sys} \quad (n = \text{number of moles; } dn=0 \text{ for closed system})$$

U is state function  $\Rightarrow dU_{sys}$  is exact differential

↓

$dn = 0$  (closed system)

$$dU(T, P) = \left(\frac{\partial U}{\partial T}\right)_P dT + \left(\frac{\partial U}{\partial P}\right)_T dP = dq_{sys} + \delta w_{sys}$$

OR

$$dU(T, V) = \left(\frac{\partial U}{\partial T}\right)_V dT + \left(\frac{\partial U}{\partial V}\right)_T dV = dq_{sys} + \delta w_{sys}$$

4

"divide through by ??"

math handout #6

$$dU(T, P) = \left(\frac{\partial U}{\partial T}\right)_P dT + \left(\frac{\partial U}{\partial P}\right)_T dP$$

"divide through by dV holding x (something else) constant"

$$\left(\frac{\partial U}{\partial V}\right)_X = \left(\frac{\partial U}{\partial T}\right)_P \left(\frac{\partial T}{\partial V}\right)_X + \left(\frac{\partial U}{\partial P}\right)_T \left(\frac{\partial P}{\partial V}\right)_X$$

later special simplification if x=P or T

5

two relationships for ideal gasses: a ( $\mu$ sec) look ahead  
*(will prove rigorously in next lecture, but gthis is the next lecture)*

- for any substance  
 $dU_V = dq_V = n\bar{C}_V dT$  and  $\Delta U_V = [n\bar{C}_V \Delta T]$  for a constant volume process
- but for an ideal gas  
 $dU = n\bar{C}_V dT$  and  $\Delta U = n\bar{C}_V \Delta T$  for ANY path (not only constant V process)  
*[other parts of path, changes of P and V with constant T, give zero contribution to  $\Delta U$ ]*

- for ideal gas  
 $\bar{C}_P = \bar{C}_V + R$
- monatomic ideal gas  
 $\bar{C}_V = \frac{3}{2}R$     $\bar{C}_P = \frac{5}{2}R$   
*[simple proof coming soon]*

6

# Chemistry 163B Winter 2014

## notes for lecture 5

**ideal gas**  $\Delta U = n\bar{C}_v\Delta T$  for ANY path (not only constant V process)

---

$dU_v = dq_v = n\bar{C}_v dT$  and  $\Delta U_v = n\bar{C}_v \Delta T$  for a constant volume process  
 $dU = n\bar{C}_v dT$  and  $\Delta U = n\bar{C}_v \Delta T$  for ANY path (not only constant V process)

(general,  $w_{other}=0, dn=0$ )

$$dU = \left(\frac{\partial U}{\partial T}\right)_V dT + \left(\frac{\partial U}{\partial V}\right)_T dV$$

$$dU = dq - PdV$$

$$dU_v = \left(\frac{\partial U}{\partial T}\right)_V dT = dq_v = n\bar{C}_v dT$$

$$\left(\frac{\partial U}{\partial T}\right)_V = n\bar{C}_v$$

$$dU = n\bar{C}_v dT + \left(\frac{\partial U}{\partial V}\right)_T dV$$

ideal gas

$$dU = n\bar{C}_v dT + \left(\frac{\partial U}{\partial V}\right)_T dV$$

but

$$\left(\frac{\partial U}{\partial V}\right)_T = 0$$

$$dU = n\bar{C}_v dT$$

$$\Delta U = n\bar{C}_v \Delta T$$

even if V not constant (i.e. any path)

$\Delta U$  along general path where both T (const V) and V (const T) vary  
 $\Delta U = n\bar{C}_v \Delta T$

7

**$C_p = C_v + nR$  for ideal gas**

---

for only P-V work and closed system ( $\delta w_{other} = 0, dn=0$ )  
 $dU = dq - P_{ext}dV$

$$dU_v = dq_v = n\bar{C}_v dT \quad dU_p = dq_p - PdV_p$$

$$dq_p = dU_p + PdV_p$$

$$\left(\frac{dq}{dT}\right)_p = \left(\frac{\partial U}{\partial T}\right)_p = n\bar{C}_v \quad \left(\frac{dq}{dT}\right)_p = n\bar{C}_p = \left(\frac{\partial U}{\partial T}\right)_p + P\left(\frac{\partial V}{\partial T}\right)_p$$

for monatomic ideal gas  
 $PV = nRT$  and  $U = \frac{3}{2}nRT$

$$\left(\frac{\partial U}{\partial T}\right)_p = \frac{3}{2}nR = \left(\frac{\partial U}{\partial T}\right)_p$$

$$n\bar{C}_p = \frac{3}{2}nR + P\left(\frac{nR}{P}\right) = \frac{5}{2}nR$$

8

**1st Law recapitulation**

---

$U \equiv$  internal energy  
 $dU_{sys} = dq_{sys} + \delta w_{sys} + dn_{sys}$  ( $n$ =number of moles;  $dn=0$  for closed system)  
 $dU_{sys} = -dU_{sur}$  (energy conserved)  
 $dU$  is exact differential  
 $U$  is a state function completely general

---

for only P-V work and closed system ( $dn=0$ )  
 $dU = dq - P_{ext}dV$

- Constant volume process  $dU_v = dq_v, \Delta U_v = q_v$
- Adiabatic process  $dU = \delta w, \Delta U = w$

9

**enthalpy:  $q$  for process at constant Pressure**

---

$H \equiv U + P_{int}V$  (definition of enthalpy, H)

since U is state function and P, V are state variables, H is also a

**STATE FUNCTION** completely general

why a new state function you might ask??

$$dU_v = dq_v; \Delta U_v = q_v \text{ heat at constant volume}$$

but most reactions and many physical processes are carried out at constant P

desire state function for  $q_p$ , heat at constant pressure

10

**enthalpy: H, a state function for heat transfer at constant pressure**

---


$$H \equiv U + P_{int}V$$

$$dH = dU + PdV + VdP$$

$$dH = dq - PdV + \delta w_{other} + PdV + VdP$$

$$dH = dq + VdP + \delta w_{other}$$

and at P=constant and  $\delta w_{other} = 0$

$$dH_p = dq_p$$

$$\Delta H_p = q_p \text{ as advertised !!}$$

$\Delta H_p = q_p$  at const P no  $w_{other}$   
 $\Delta H_p > 0$  endothermic (heat gained by system)  
 $\Delta H_p < 0$  exothermic (heat lost by system)

11

**$\Delta H$  ideal gas**

---


$$\Delta H_p = q_p = \int n\bar{C}_p dT \approx n\bar{C}_p \Delta T \text{ (general, } w_{other}=0, dn=0)$$

ideal gas

$$H \equiv U + PV = U + nRT$$

$$dH = dU + nRdT \text{ (general for ideal gas)}$$

$$dH = n\bar{C}_v dT + nRdT \text{ (general for ideal gas, even V not const)}$$

$$dH = n(\bar{C}_v + R)dT$$

$$dH = n\bar{C}_p dT \text{ IDEAL GAS ANYTIME, EVEN IF P NOT CONSTANT}$$

$$\Delta H = n\bar{C}_p \Delta T \text{ ideal gas general (} w_{other}=0, dn=0)$$

12

# Chemistry 163B Winter 2014

## notes for lecture 5

### manipulating thermodynamic functions: fun and games

for example:  
HW#3

12. Derive the following for any closed system, with only P-V work:

$$C_V = - \left( \frac{\partial U}{\partial V} \right)_T \left( \frac{\partial V}{\partial T} \right)_U$$

13

### total differential for $U(T,V,n)$ and $H(T,P,n)$

$$U(T, V, n_1, n_2, \dots, n_N)$$

$$dU = \left( \frac{\partial U}{\partial T} \right)_{V,n} dT + \left( \frac{\partial U}{\partial V} \right)_{T,n} dV + \sum_{i=1}^N \left( \frac{\partial U}{\partial n_i} \right)_{T,V,n_{j \neq i}} dn_i$$

$$H(T, P, n_1, n_2, \dots, n_N)$$

$$dH = \left( \frac{\partial H}{\partial T} \right)_{P,n} dT + \left( \frac{\partial H}{\partial P} \right)_{T,n} dP + \sum_{i=1}^N \left( \frac{\partial H}{\partial n_i} \right)_{T,P,n_{j \neq i}} dn_i$$

for now closed system all  $dn_i = 0$

14

### $H(T,P)$ : some manipulations and relationships (closed system)

$$dH = \bar{n}dq + VdP \quad \text{closed system, } \bar{n}v_{\text{other}} = 0$$

'divide by dT, holding P constant'

$$\left( \frac{\partial H}{\partial T} \right)_P = \left( \frac{dq}{dT} \right)_P + V \left( \frac{\partial P}{\partial T} \right)_P$$

$$\left( \frac{\partial H}{\partial T} \right)_P = \left( \frac{dq}{dT} \right)_P = n\bar{c}_P$$

'divide by dP, holding T constant'

$$\left( \frac{\partial H}{\partial P} \right)_T = \left( \frac{dq}{dP} \right)_T + V \left( \frac{\partial P}{\partial P} \right)_T$$

$$\left( \frac{\partial H}{\partial P} \right)_T = \left( \frac{dq}{dP} \right)_T + V$$

$$dH = \left( \frac{\partial H}{\partial T} \right)_P dT + \left( \frac{\partial H}{\partial P} \right)_T dP$$

$$dH = n\bar{c}_P dT + \left( \frac{\partial H}{\partial P} \right)_T dP \quad \text{eqns. 3.30-3.32 E&R (p. 56 [§2]_...)} \quad \text{math handout \#6}$$

15

### $U(T,V)$ : some manipulations and relationships (closed system)

$$dU = \bar{n}dq - PdV \quad \text{closed system, } \bar{n}v_{\text{other}} = 0$$

'divide by dT, holding V constant'

$$\left( \frac{\partial U}{\partial T} \right)_V = \left( \frac{dq}{dT} \right)_V - P \left( \frac{\partial V}{\partial T} \right)_V$$

$$\left( \frac{\partial U}{\partial T} \right)_V = \left( \frac{dq}{dT} \right)_V = n\bar{c}_V$$

'divide by dV, holding T constant'

$$\left( \frac{\partial U}{\partial V} \right)_T = \left( \frac{dq}{dV} \right)_T - P \left( \frac{\partial V}{\partial V} \right)_T$$

$$\left( \frac{\partial U}{\partial V} \right)_T = \left( \frac{dq}{dV} \right)_T - P$$

$$dU = \left( \frac{\partial U}{\partial T} \right)_V dT + \left( \frac{\partial U}{\partial V} \right)_T dV$$

$$dU = n\bar{c}_V dT + \left( \frac{\partial U}{\partial V} \right)_T dV \quad \text{eqn. 3.12-3.15 E&R (p 50 [§6]_...)} \quad \text{math handout \#6}$$

16

### save for later when we have tools from 2<sup>nd</sup> Law of Thermodynamics

$$dU = n\bar{c}_V dT + \left( \frac{\partial U}{\partial V} \right)_T dV \quad \text{need 2<sup>nd</sup> Law to evaluate this in terms of P,V,T}$$

$$\left( \frac{\partial U}{\partial V} \right)_T = T \left( \frac{\partial P}{\partial T} \right)_V - P \quad \text{E \& R eqn. 3.19}$$

many of the results in E&R ch 3 use this [yet] 'unproven' result; we will derive later class should use result in HW3 #13\*

17

### some important relationships between $C_P$ and $C_V$

$$\frac{1}{n} \left( \frac{\partial H}{\partial T} \right)_P = \bar{c}_P \quad \text{and} \quad \frac{1}{n} \left( \frac{\partial U}{\partial T} \right)_V = \bar{c}_V$$

to get relationship between  $C_P$  and  $C_V$  one needs to have relationship involving both H and U; soooo

$$H = U + PV$$

$$dH = dU + PdV + VdP$$

18

# Chemistry 163B Winter 2014

## notes for lecture 5

**continuing with relating  $C_p$  and  $C_v$**

---


$$H = U + PV$$

$$dH = dU + PdV + VdP$$

divide by  $dT$ ,  $P$  constant

$$\left(\frac{\partial H}{\partial T}\right)_P = \left(\frac{\partial U}{\partial T}\right)_P + P\left(\frac{\partial V}{\partial T}\right)_P + V\left(\frac{\partial P}{\partial T}\right)_P$$

$$n\bar{C}_p = \left(\frac{\partial U}{\partial T}\right)_P + P\left(\frac{\partial V}{\partial T}\right)_P$$

now to get  $\left(\frac{\partial U}{\partial T}\right)_P$

$$dU = n\bar{C}_v dT + \left(\frac{\partial U}{\partial V}\right)_T dV$$

$$\left(\frac{\partial U}{\partial T}\right)_P = n\bar{C}_v + \left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_P$$

19

**let's finish  $C_p$  vs  $C_v$  (very general relationship) -E&R 3.37**

---


$$n\bar{C}_p = \left(\frac{\partial U}{\partial T}\right)_P + P\left(\frac{\partial V}{\partial T}\right)_P$$

$$n\bar{C}_p = n\bar{C}_v + \left(\frac{\partial U}{\partial V}\right)_T \left(\frac{\partial V}{\partial T}\right)_P + P\left(\frac{\partial V}{\partial T}\right)_P$$

$$n\bar{C}_p = n\bar{C}_v + \left[\left(\frac{\partial U}{\partial V}\right)_T + P\right] \left(\frac{\partial V}{\partial T}\right)_P$$

Annotations:

- energy to raise T 1° const P (vol changes) →  $\left(\frac{\partial U}{\partial T}\right)_P$
- energy to raise T 1° const V →  $n\bar{C}_v$
- potential energy as molecules 'separate' per unit volume change →  $\left(\frac{\partial U}{\partial V}\right)_T$
- energy lost as P-V work per unit volume change →  $P$
- volume change per T change of 1° →  $\left(\frac{\partial V}{\partial T}\right)_P$

20

**$C_p$  vs  $C_v$  for ideal gas**

---


$$n\bar{C}_p = n\bar{C}_v + \left[\left(\frac{\partial U}{\partial V}\right)_T + P\right] \left(\frac{\partial V}{\partial T}\right)_P$$

for ideal gas

$$V = \frac{nRT}{P}$$

Energy, U is function of ONLY T, U(T)

$$\left(\frac{\partial U}{\partial T}\right)_P = \frac{nR}{P}$$

$$\left(\frac{\partial U}{\partial V}\right)_T = 0$$

$$n\bar{C}_p = n\bar{C}_v + [0 + P] \frac{nR}{P}$$

$$n\bar{C}_p = n\bar{C}_v + nR$$

$\bar{C}_p = \bar{C}_v + R$  for ideal gas

21

**experimental  $C_v$  and  $C_p$  for selected gases**

---

Nature of gas	Gas	$C_v$ (J mol <sup>-1</sup> K <sup>-1</sup> )	$C_p$ (J mol <sup>-1</sup> K <sup>-1</sup> )	$C_p - C_v$ (J mol <sup>-1</sup> K <sup>-1</sup> )	$\gamma$
Monatomic	He	12.5	20.8	8.30	1.66
Monatomic	Ne	12.7	20.8	8.12	1.64
Monatomic	Ar	12.5	20.8	8.30	1.67
Diatomic	H <sub>2</sub>	20.4	28.8	8.45	1.41
Diatomic	O <sub>2</sub>	21.0	29.3	8.32	1.40
Diatomic	N <sub>2</sub>	20.8	29.1	8.32	1.40
Triatomic	H <sub>2</sub> O	27.0	35.4	8.35	1.31
Polyatomic	CH <sub>4</sub>	27.1	35.4	8.36	1.31

$R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$ 

ideal gas

$$\bar{C}_p - \bar{C}_v = R$$

monatomic  $\bar{C}_v = \frac{3}{2}R$

diatomic  $\bar{C}_v \approx \frac{5}{2}R$

$\frac{3}{2}R = 12.47$

$\frac{5}{2}R = 20.78$

$\frac{7}{2}R = 29.10$

Table from: <http://www.scribd.com/doc/33638936/NCERT-Book-Physics-Class-XI-2>

22

**in section derive equation following equation**

---


$$n\bar{C}_v = n\bar{C}_p + \left[\left(\frac{\partial H}{\partial P}\right)_T - V\right] \left(\frac{\partial P}{\partial T}\right)_V$$

start with

$$dU = dH - PdV - VdP$$

divide by  $dT$  with  $V$  constant

and then boogie along as we just did!!

23

**First Law: ideal gas calculations**

---

relationships that apply to ideal gasses for all conditions with  $w_{\text{other}}=0$  and constant composition (some also apply more generally):

$\Delta U = q + w$	$w_{P_{\text{ext}}} = -\int P_{\text{ext}} dV$	$PV = nRT$
$q_V = n\int \bar{C}_v dT$ ? $= n\bar{C}_v \Delta T$	$q_P = n\int \bar{C}_p dT$ ? $= n\bar{C}_p \Delta T$	$\bar{C}_p = \bar{C}_v + R$
$H \equiv U + PV$	$\Delta U_{\text{any conditions}} = n\bar{C}_v \Delta T$	$\Delta H_{\text{any conditions}} = n\bar{C}_p \Delta T$
monatomic ideal gas	$\bar{C}_v = \frac{3}{2}R$	$\bar{C}_p = \frac{5}{2}R$

24