

Chemistry 163B, Winter 2020  
Lecture 1- Introductory Lecture


Chemistry 163B  
Thermodynamics  
Winter 2020

1

1

*Menu: for TODAY*

---



- Introduction and Class Resources, etc.
- Careful definition of thermodynamic variables ( $P, V, T$ , etc)
- Ideal gas ENERGY DEPENDS ONLY ON TEMPERATURE (*derivation*)
- Van der Waals equation (*example of equation of state for 'real' gas*)

2

2

Thermodynamics is a  
really beautiful scientific  
story !!

3

3

*"rules" of science*

---

- observations
- ↓
- guiding principles
- ↓
- predictions and applications based on principles

4


4

*Observations (QM vs Thermo)*

---

Thermodynamics is very 'working class' in its origins:


quantum mechanics



**Effete**

$H \Psi = E \Psi$

thermodynamics



**Blue Collar**

$\Delta S_{\text{universe}} > 0$

5

5

*observations: thermo = heat*

---

- Count Rumford, 1799
- observed water turning into steam when canon barrel was bored
- $work \leftrightarrow heat$






FIGURE 11. An illustration from Rumford's paper, "An Inquiry Concerning the Source of the Heat Which is Excited by Friction," showing the apparatus used by him in the cannon boring experiment. Figure 1, upper left, shows the cannon as received from the foundry, and Figure 2, below, shows it mounted in the machine used for boring. (Reproduced with the permission of Harvard University Press.)

6




6

# Chemistry 163B, Winter 2020

## Lecture 1- Introductory Lecture

observations: mechanical efficiency of steam engine




- Sadi Carnot, 1824
- efficiency of engines

7

guiding principles

- Conservation of heat and work (Joule, 1845)  
**1st LAW OF THERMODYNAMICS**
- Clausius, 1860 Entropy  
**2nd LAW OF THERMODYNAMICS**
- Boltzmann, late 19th century,  
**molecular picture of entropy**

Clausius  
Boltzmann

the thermodynamic functions **U, H, and S**  
(1st and 2nd laws)

8


7

8

"Applications" **U,H,S,G**

How does knowledge about efficiencies of steam engines, mechanical systems, etc. relate to processes in chemical, biological, and geological systems?

ANSWERED BY:




J. W. Gibbs- arguably the first great American scientist who combined the concepts of heat and entropy and proposed "[Gibbs] Free Energy", **G**, a thermodynamic state function that leads to a whole spectrum of applications

9

9

from: Einstein's "Autobiographical Notes"



A theory is the more impressive the greater the simplicity of its premises, the more different kinds of things it relates, and the more extended its area of applicability. Hence the deep impression that **classical thermodynamics** made upon me. It is the only physical theory of universal content concerning which I am convinced that, within the framework of the applicability of its basic concepts, **it will never be overthrown** (for the special attention of those who are skeptics on principle).

10


10

types of problems that thermodynamics addresses (reactions)

**might**

1. Under what conditions ~~will~~ a reaction occur ?

EXAMPLE:  
 $C(\text{graphite}) \rightarrow C(\text{diamond})$  1 atm 398K **nada**  
 30,000 -100,000 atm **jHurra!**  
 1000 K - 3000 K  
 1st artificially produced diamonds, 1954 at General Electric Labs



11


11

types of problems that thermodynamics addresses (equilibria)


2. How far will a reaction proceed (given enough time)?  
 How do the thermal and entropic properties determine EQUILIBRIUM ?  
 How do the EQUILIBRIUM conditions depend on T,P ?

$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$

what are best T,P for  $NH_3$  products?  
**The Haber Process**



$SiO_2 + CaCO_3 \rightleftharpoons CaO + SiO_2 + CO_2(g)$   
 quartz calcite wollastonite



**geologic thermometer.**  
 The fraction of wollastonite in a rock sample can be used to estimate the temperature at which the rock metamorphosis occurred [the T at which the equilibrium was rapidly frozen]

**Geologic Thermometer**

12

12

# Chemistry 163B, Winter 2020

## Lecture 1- Introductory Lecture

*types of problems that thermodynamics addresses (biology)*


3. Chemical and physical changes in biological systems

$[K^+]_{\text{inside}}$

$\rightleftharpoons$

$[K^+]_{\text{outside}}$

cell      what difference in  $[K^+]_{\text{out}}$  vs  $[K^+]_{\text{in}}$  can be tolerated before cell wall bursts or collapses?




low hypotonic (rupture)
normal isotonic
high hypertonic (collapse)

<http://www.vivo.colostate.edu/hbooks/cmb/cells/pmembr/osmosis.html>

13


*types of problems that thermodynamics addresses (biology)*

4. Why does an egg hard boil? (protein conformation)



Uncooked Proteins      Cooked proteins network

5. Membrane potentials and ion concentrations in neurons. (electrochemistry and thermodynamics)




Cytosol      Extracellular Membrane

Distribution of electrical charge across the membrane. The uneven charges inside and outside the neuron line up along the membrane because of electrostatic attraction across this very thin barrier.

14

*types of problems that thermodynamics addresses (ecology)*

6. Thermodynamic feasibility of  $SO_2$  removal

$$SO_2(g) \rightleftharpoons S(s) + O_2(g)$$


<http://healthandenergy.com/images/magnifica%20smoke%20stacks.jpg>

15

*what thermodynamics can't answer*

- How fast a reaction proceeds (kinetics, catalysts, enzymes; in chem 103, BMB 100, chem 163C)
- Macroscopic thermodynamics does not prove or require hypotheses about molecular structure; however we will use our knowledge of molecular structure to get an atomic "picture" of thermodynamic processes. The quantitative connection is made by statistical thermodynamics.
 

$\text{chem 163A} \rightleftharpoons \text{chem 163B}$   
 $\text{chem 163C}$
- Although in chemistry 163B we will study how thermodynamics put limits on processes at equilibrium, there exists a whole other field of *non-equilibrium thermodynamics*.

16

*aims of class*

- Clear conceptual picture of thermodynamics
- Ability to relate and apply thermodynamics to chemical and biological systems
- PROBLEM SOLVING: Chemistry + Logic + Mathematics**
- How to do independent and advanced reading/research in areas that utilize thermodynamics.
- THE GRAND PICTURE of how thermodynamics and quantum mechanics fit into our picture of 'nature'.
- Advanced mathematical techniques


17

*class administration*

[www.chemistry.ucsc.edu](http://www.chemistry.ucsc.edu)  $\Rightarrow$  course pages  $\Rightarrow$  Chemistry 163B

<http://switkes.chemistry.ucsc.edu/teaching/CHEM163B/>

- lectures: **A MUST**
- homework: **A MUST**
- sections:        required (**A MUST**)
- tutorial EVENT & Office Hours & LSS (for YOU!)
- midterms: 31<sup>st</sup> January  
28<sup>th</sup> February
- final: 19<sup>th</sup> March, 4:00-7:00 PM  
(last class 13<sup>th</sup> March)




18

# Chemistry 163B, Winter 2020

## Lecture 1- Introductory Lecture

### Learn Thermodynamics

- **Lectures**
- **Sections** (start Monday, Jan 13; HW#1 Probs 1-8)
- **Tutorial Event** (starts TOMORROW Tues, Jan 7, HW#1 Probs1-2)
- **Office hours** (start TODAY Mon, Jan 6)
- **LSS Tutor**



19

19

**first Tutorial Tuesday 7<sup>th</sup> January, NS Annex 102 , 5:25PM-6:30PM**

---

Chemistry 163B Winter 2020 Chemistry 16

**Comments on Mathematical Techniques for Chemistry 163B**

Our text (Zumdahl, 8E, 2017) pp. 62-63, 65-68, 70-71, 73, 75, contains detailed information on the topics of differential and integral calculus relevant to the study of thermodynamics. It is not meant to be a replacement for the usual calculus textbooks and homework problems, rather than just as detailed mathematical exercises. However, it is useful for conceptualizing the topics. In your first discussion section the TA will also comment on the techniques.

1. Partial derivatives:
  - a. Know the physical meaning of partial derivatives.
  - b. Know how to calculate them mechanically.
  - c. Know chain rule.
2. You should have "a hand" for the derivatives and integrals of functions like  $e^{ax}$ ,  $\ln(x)$ ,  $e^{ax} \sin bx$ ,  $\sin ax$ . (I'll assist you in your necessary memory.)
3. Differential expressions (see 2.6, pp 39-41, 63) are in  $dx$  form:
  - a.  $\int dx$  is exact differential if  $\int dx$  depends on the path from initial  $\rightarrow$  final. This implies there is no "underlying" well behaved function  $y$  that changes with  $x$ .
  - b.  $\int dx$  is exact differential if  $\int dx$  does not depend on the path from initial  $\rightarrow$  final.
4. Implications that arise from  $\int dx$  being an exact differential:
  - a. There is a well behaved function  $y$  whose change is described by  $\int dx$ .
  - b.  $\int dx = \int \left( \frac{\partial y}{\partial x} dx + \frac{\partial y}{\partial z} dz \right)$  as  $y$  is a many general variable.
  - c.  $\oint dx = 0$  (integral for any cyclic path, i.e. the initial and final states are identical).
  - d.  $\int dx = y(\text{final}) - y(\text{initial})$ .

20

20

Winter 2020 Chemistry 163B

**Tutorial Tuesday 7<sup>th</sup> January**

**Homework Problems (#1-#8)**

1. For an ideal gas  $P\bar{V} = RT$  ( $\bar{V} \equiv V_m \equiv \frac{V}{n}$  molar volume) evaluate:
  - a.  $\left( \frac{\partial P}{\partial \bar{V}} \right)_T$
  - b.  $\left( \frac{\partial \bar{V}}{\partial T} \right)_P$
  - c.  $\left( \frac{\partial T}{\partial P} \right)_P$
  - d.  $\left( \frac{\partial P}{\partial \bar{V}} \right)_T \left( \frac{\partial \bar{V}}{\partial T} \right)_P \left( \frac{\partial T}{\partial P} \right)_P$
2. For a Van der Waals gas:  $\left( P + \frac{a}{\bar{V}^2} \right) (\bar{V} - b) = RT$  evaluate:
  - a.  $\left( \frac{\partial P}{\partial \bar{V}} \right)_T$
  - b.  $\left( \frac{\partial^2 P}{\partial \bar{V}^2} \right)_T$
  - c.  $\left( \frac{\partial P}{\partial T} \right)_P$
  - d.  $\left[ \frac{\partial}{\partial T} \left( \frac{\partial P}{\partial \bar{V}} \right)_T \right]_P$
  - e.  $\left[ \frac{\partial}{\partial \bar{V}} \left( \frac{\partial P}{\partial T} \right)_P \right]_T$

21

21

Winter 2020 Chemistry 163B

3. Engel & Reid problem P1.38 (4<sup>th</sup> edition)

to HW#1 listing on CANVAS Van der Waals gas

4. (optional) The Van der Waals and virial expressions are two commonly used equations of state as approximations to real gas behavior:

$$P = \frac{RT}{(\bar{V} - b)} - \frac{a}{\bar{V}^2} \quad \text{Van der Waals}$$

$$P = RT \left[ \frac{C_0(T)}{\bar{V}} + \frac{C_1(T)}{\bar{V}^2} + \frac{C_2(T)}{\bar{V}^3} + \frac{C_3(T)}{\bar{V}^4} \right] \quad \text{virial}$$

- a. Show that the first three virial coefficients are related to the Van der Waals parameters  $a$  and  $b$  in the following way:
  - $C_0(T) = 1$
  - $C_1(T) = b - a/RT$
  - $C_2(T) = b^2$

*HINT:* note that the virial expansion is just a Taylor (Maclaurin) series in  $x = \left( \frac{b}{\bar{V}} \right)$

$$\frac{P}{RT} = \sum_{i=0}^{\infty} C_i(T) \left( \frac{b}{\bar{V}} \right)^i = \sum_{i=0}^{\infty} C_i(T) x^i = f(x; T)$$

with coefficient  $C_i(T) = \frac{1}{i!} \left( \frac{d^i f}{dx^i} \right)_{x=0}$
- b. Why does the Van der Waals "a" only appear in  $C_1(T)$ ?

22

22

enough hellos ....

and NOW TO BEGIN !!


our thermodynamics trek





23

23


system and surroundings (careful definitions)  E&R<sub>4th</sub> sec 1.3

A thermodynamic **system** consists of all the materials involved in the process under study. This material could be the contents of an open beaker containing reagents, the electrolyte solution within an electrochemical cell, or the contents of a cylinder and movable piston assembly in an engine.

In thermodynamics, the rest of the universe is referred to as the **surroundings**.

system + surroundings = the **UNIVERSE**

If a system can exchange matter with the surroundings, it is called an **open system**; if not, it is a **closed system**.



24

24



# Chemistry 163B, Winter 2020

## Lecture 1- Introductory Lecture

*heuristic derivation*

8.  $\frac{n^*}{V}$  is density of molecules.  $Av_x\Delta t = V$  is volume of rectangular box,

9.  $\frac{1}{2} \frac{n^*}{V} Av_x\Delta t$  is total number of molecules colliding with area A in  $\Delta t$

10. total  $\Delta p = (2mv_x) \left( \frac{1}{2} \right) \left( \frac{n^*}{V} \right) (Av_x\Delta t) = (mv_x^2) \left( \frac{n^*}{V} \right) (A\Delta t)$

31

31

*heuristic derivation*

11. after some algebra and equating

$$P = \frac{F}{A} = \frac{\Delta p}{A\Delta t} = (mv_x^2) \left( \frac{n^*}{V} \right) = \frac{n^*kT}{V}$$

$$P = \left( \frac{mv_x^2}{2} \right) \left( \frac{n^*}{V} \right) = \left( \frac{v_x^2}{V} \right) \frac{kT}{2}$$

$\left( \frac{mv_x^2}{2} \right) = \frac{kT}{2}$  for each molecule of 1-D ideal gas

12. in 3D with  $|v_x| = |v_y| = |v_z|$

$$\text{total KE} = \left( \frac{1}{2}mv_x^2 + \frac{1}{2}mv_y^2 + \frac{1}{2}mv_z^2 \right) n^* = \frac{3}{2}n^*kT = \frac{3}{2}nRT$$

$\frac{1}{2}RT$  for each deg. free (x,y,z) of 'classical freedom'

32

32

*TAKE HOME MESSAGES*

- Good warm up of physics and equation derivation
- For a molecule with only kinetic energy (e.g. monatomic species), and ideal gas (no intermolecular forces)

**$E = \frac{3}{2}kT$  per molecule or  $E = \frac{3}{2}RT$  per mole**

- For monatomic ideal gas, E is function of **only T**;

**T constant  $\leftrightarrow$  E constant**

33

33

*beyond PV=nRT: 'real' gases and equations of state (E&R sec 1.5)*

Ideal  
Prof S ??

vs

Real  
Prof S ✓✓

34

34

*various equations of state (Raff Table 1.2)  $\frac{V}{n} \equiv \bar{V} \equiv V_m$  molar volume*

Equation of State	Functional Form	Number of Parameters
Ideal gas	$PV_m = RT$	0
→ van der Waals	$(V_m - b)(P + a/V_m^2) = RT$	2
Dieterici	$P(V_m - b)\exp(a/RTV_m) = RT$	2
Berthelot	$(V_m - b)(P + a/TV_m^2) = RT$	2
→ Virial	$P = RT \left[ V_m^{-1} + \sum_{n=2}^{\infty} C_n(T)V_m^{-n} \right]$	∞
Beattie-Bridgman	$PV_m^2 = (1 - \gamma)RT(V_m + \beta) - \alpha$ with $\gamma = c_v/T^2V_m$ $\beta = b_1(1 - b/V_m)$ , and $\alpha = a_1(1 + a/V_m)$	5
Redlich-Kwong	$P = \frac{RT}{(V_m - b)} - \frac{a}{T^{1/2}V_m(V_m + b)}$	2
Reichsanstalt	$PV = RT + AP + BP^2 + CP^3 + \dots$	∞

35

35

Some comments on the  
Van der Waals equation of state

36

36

**Slide 32**

---

**ES2**

Eugene Switkes, 1/6/2020


# Chemistry 163B, Winter 2020

## Lecture 1- Introductory Lecture

### van der Waals equation the parameter b

ideal gas law  $P_{ideal} \bar{V}_{ideal} = RT$

van der Waals  $\left(P + \frac{a}{\bar{V}^2}\right) (\bar{V} - b) = RT$



interpretation of parameters:

- b is correction for actual volume of atoms/molecules
  - volume available to molecules  $(\bar{V} - b)$  smaller than  $\bar{V}$
  - b is associated with repulsive forces


37

37

### van der Waals equation the parameter a

$P_{ideal} \bar{V}_{ideal} = RT$

$\left(P + \frac{a}{\bar{V}^2}\right) (\bar{V} - b) = RT$



interpretation of parameters:



- a is correction for attractive forces of atoms/molecules
  - a is associated with attractive forces
  - $P_{bulk} \equiv P_{ideal}$  is greater than  $P_{meas}$  measured at surface
  - $P_{bulk} = \left(P_{meas} + \frac{a}{\bar{V}^2}\right)$   $P_{meas} \rightarrow P_{bulk}$   $\bar{V} \rightarrow \infty$

38

38

### heuristic justification for attractive constant a

- asymmetric attractive forces for molecule at surface
- molecule at surface has less momentum less than molecule in bulk
- $P_{meas} < P_{bulk}$
- $P_{bulk} = \left(P + \frac{a}{\bar{V}^2}\right)$

39

39

### van der Waals equation

$$\left(P + \frac{a}{\bar{V}^2}\right) (\bar{V} - b) = RT$$

Gas	$b$ (L mol <sup>-1</sup> )	$a$ (L <sup>2</sup> bar mol <sup>-2</sup> )
He	0.0238	0.0346
Ne	0.01672	0.208
Ar	0.03201	1.355
Kr	0.0396	2.325
Acetylene	0.0522	4.516
N <sub>2</sub>	0.0387	1.37
H <sub>2</sub> O	0.03049	5.537
CO <sub>2</sub>	0.04286	3.658

Source: Handbook of Chemistry and Physics, 78th edition, CRC Press, Boca Raton, FL, 1997-98

40

40

### van der Waals equation

$$\left(P + \frac{a}{\bar{V}^2}\right) (\bar{V} - b) = RT$$

Gas	$b$ (L mol <sup>-1</sup> )	$a$ (L <sup>2</sup> bar mol <sup>-2</sup> )
He	0.0238	0.0346
Ne	0.01672	0.208
Ar	0.03201	1.355
Kr	0.0396	2.325
Acetylene	0.0522	4.516
N <sub>2</sub>	0.0387	1.37
H <sub>2</sub> O	0.03049	5.537
CO <sub>2</sub>	0.04286	3.658

Source: Handbook of Chemistry and Physics, 78th edition, CRC Press, Boca Raton, FL, 1997-98

41

41

some trash talk on VDW eqn: **inaccurate but instructive !!!** (from other texts)

**Validity**

However, the Van der Waals model is not appropriate for rigorous quantitative calculations, remaining useful only for **teaching** and qualitative purposes.<sup>[1]</sup>

Nowadays, Eq. 2.9 belongs to **“pedagogical physics:”** it is the simplest equation that illustrates several important concepts, but its accuracy is not satisfactory.

42


42



# Chemistry 163B, Winter 2020

## Lecture 1- Introductory Lecture

Menu: for TODAY



- ✓ Introduction and Class Resources, etc.
- ✓ Careful definition of thermodynamic variables ( $P$ ,  $V$ ,  $T$ , etc)
- ✓ Ideal gas ENERGY DEPENDS ONLY ON TEMPERATURE ( $T$ )
- ✓ Van der Waals equation (example of equation of state for 'real' gas)

43

43

## End of Lecture 1

44

44

various equations of state (Raff Table 1.2)

Equation of State	Functional Form	Number of Parameters
Ideal gas	$PV_m = RT$	0
van der Waals	$(V_m - b)(P + a/V_m^2) = RT$	2
Dieterici	$P(V_m - b)\exp[a/RTV_m] = RT$	2
Berthelot	$(V_m - b)(P + a/TV_m^2) = RT$	2
Virial	$P = RT \left[ V_m^{-1} + \sum_{n=2}^{\infty} C_n(T) V_m^{-n} \right]$	$\infty$
Beattie-Bridgman	$PV_m^2 = (1 - \gamma)RT(V_m + \beta) - \alpha$ with $\gamma = c_0/T^2 V_m$ , $\beta = b_0(1 - b/V_m)$ , and $\alpha = a_0(1 + a/V_m)$	5
Redlich-Kwong	$P = \frac{RT}{(V_m - b)} - \frac{a}{T^{1/2} V_m(V_m + b)}$	2
Reichsanstalt	$PV = RT + AP + BP^2 + CP^3 + \dots$	$\infty$

45

45

### The Haber Process (thermodynamics and kinetics)

Previously the problem had been that  $N_2$  is a very stable molecule, and so most attempts to convert it to less stable molecules, such as  $NH_3$ , failed because of **thermodynamic or entropy problems**. The secret to the Haber-Bosch process proved to be a **catalyst of iron** with a small amount of **aluminium** added (aluminium was at the time an exotic and expensive metal that probably attracted Haber's attention as a novelty). The Haber-Bosch process operates at **high pressure** so as to shift the equilibrium to the right, and **high temperature to increase the rates of the reaction**. Of course, operating at high temperature actually shifted the reaction to the left, but the trade-off for faster rates was accepted. By removing the ammonia as liquid ammonia, the equilibrium is continuously shifted to the right.


46

46

### heuristic

heu-ris-tic [*h*voo-ris-tik or, often, yoo-] adjective


1. serving to indicate or point out; stimulating interest as a means of furthering investigation.



47

47


### universal behavior of gas



48

48

# Chemistry 163B, Winter 2020 Lecture 1- Introductory Lecture



49