

Lecture 15 Chemistry 163B
Free Energy
and
Equilibrium
E&R (\approx ch 6)



1

$\Delta G_{\text{reaction}}$ and equilibrium (first pass, lecture 14)

here $\Delta G \Rightarrow \Delta G_{\text{reaction}}$

1. $\Delta G < 0$ spontaneous ('natural', irreversible)
 $\Delta G = 0$ equilibrium (reversible)
 $\Delta G > 0$ spontaneous in reverse direction
2. $\Delta G_T = \Delta H - T\Delta S$
3. ΔG° all reactants and products in standard states
4. $\Delta \bar{G}_f^\circ \equiv \bar{G}_f^\circ$ Appendix A at 298.15K (reaction where reactants are elements in their most stable form and in their standard states, $P=1$ atm, $[\text{conc}]=1M$, etc)
 $\Delta \bar{G}_f^\circ (O_2(g)) \equiv 0$ $\Delta \bar{G}_f^\circ (C(\text{gr})) = 0$

5.

$$\Delta G_{\text{reaction}}^\circ = \sum_i v_i \Delta \bar{H}_f^\circ - T \sum_i v_i \bar{S}_i^\circ$$
$$\Delta G_{\text{reaction}}^\circ = \Delta H_{\text{reaction}}^\circ - T \Delta S_{\text{reaction}}^\circ$$

NOTE : in Appendix A: $\Delta \bar{G}_f^\circ$ and $\Delta \bar{H}_f^\circ$ in kJ mol^{-1} BUT \bar{S}° in $\text{J K}^{-1} \text{mol}^{-1}$

2

goals of lecture 15

6. Brief hello to thermodynamics of multicomponent systems (n_i 's vary)

7. $\Delta G_{\text{reaction}}$ for non-standard state concentrations, pressures

$$\Delta G_{\text{reaction}} = \Delta G^{\circ}_{\text{reaction}} + RT \ln Q$$


8. K_{eq} and $\Delta G^{\circ}_{\text{reaction}}$

9. $\Delta G_{\text{reaction}} = \Delta G^{\circ}_{\text{reaction}} + RT \ln Q$ is extensive

10. Variation of K_{eq} with T

3

6.0 molar free energy and partial molar free energy (chemical potential)



multicomponent mixture

n_{NO_2} moles NO_2 ;

$n_{\text{N}_2\text{O}_4}$ moles N_2O_4

$$dG = \left(\frac{\partial G}{\partial T} \right)_{P, n_{\text{NO}_2}, n_{\text{N}_2\text{O}_4}} dT + \left(\frac{\partial G}{\partial P} \right)_{T, n_{\text{NO}_2}, n_{\text{N}_2\text{O}_4}} dP + \left(\frac{\partial G}{\partial n_{\text{NO}_2}} \right)_{T, P, n_{\text{N}_2\text{O}_4}} dn_{\text{NO}_2} + \left(\frac{\partial G}{\partial n_{\text{N}_2\text{O}_4}} \right)_{T, P, n_{\text{NO}_2}} dn_{\text{N}_2\text{O}_4}$$

$$\mu_{\text{NO}_2} = \left(\frac{\partial G_{(\text{NO}_2 + \text{N}_2\text{O}_4 \text{ mixture})}}{\partial n_{\text{NO}_2}} \right)_{T, P, n_{\text{N}_2\text{O}_4}}$$

partial molar Gibbs free energy
or **chemical potential** (of NO_2)


more generally $G_{\text{mixture}}(T, P, n_1, \dots, n_N)$

$$dG = \left(\frac{\partial G}{\partial T} \right)_{P, n_i} dT + \left(\frac{\partial G}{\partial P} \right)_{T, n_i} dP + \sum_{i=1}^N \left(\frac{\partial G}{\partial n_i} \right)_{T, P, n_{j \neq i}} dn_i = -SdT - VdP + \sum_{i=1}^N \mu_i dn_i$$

4

Chemistry 163B Winter 2020
Lecture 15- Free Energy and Equilibrium

6.0 molar free energy and partial molar free energy (chemical potential)



multicomponent mixture

n_{NO_2} moles NO_2 ;

$n_{\text{N}_2\text{O}_4}$ moles N_2O_4

$$G_{\text{NO}_2+\text{N}_2\text{O}_4 \text{ mixture}}(T, P, n_{\text{NO}_2}, n_{\text{N}_2\text{O}_4})$$

$$dG = \left(\frac{\partial G}{\partial T}\right)_{P, n_{\text{NO}_2}, n_{\text{N}_2\text{O}_4}} dT + \left(\frac{\partial G}{\partial P}\right)_{T, n_{\text{NO}_2}, n_{\text{N}_2\text{O}_4}} dP + \left(\frac{\partial G}{\partial n_{\text{NO}_2}}\right)_{T, P, n_{\text{N}_2\text{O}_4}} dn_{\text{NO}_2} + \left(\frac{\partial G}{\partial n_{\text{N}_2\text{O}_4}}\right)_{T, P, n_{\text{NO}_2}} dn_{\text{N}_2\text{O}_4}$$


$$\mu_{\text{NO}_2} = \left(\frac{\partial G_{(\text{NO}_2+\text{N}_2\text{O}_4 \text{ mixture})}}{\partial n_{\text{NO}_2}}\right)_{T, P, n_{\text{N}_2\text{O}_4}} \quad \text{partial molar Gibbs free energy or chemical potential (of NO}_2\text{)}$$

more generally $G_{\text{mixture}}(T, P, n_1, \dots, n_N)$

$$dG = \left(\frac{\partial G}{\partial T}\right)_{P, n_i} dT + \left(\frac{\partial G}{\partial P}\right)_{T, n_i} dP + \sum_{i=1}^N \left(\frac{\partial G}{\partial n_i}\right)_{T, P, n_{j \neq i}} dn_i = -SdT - VdP + \sum_{i=1}^N \mu_i dn_i$$

5

6.1 molar free energy and partial molar free energy (chemical potential)



multicomponent mixture

n_{NO_2} moles NO_2 ;

$n_{\text{N}_2\text{O}_4}$ moles N_2O_4

$$\mu_{\text{NO}_2} = \left(\frac{\partial G_{(\text{NO}_2+\text{N}_2\text{O}_4 \text{ mixture})}}{\partial n_{\text{NO}_2}}\right)_{T, P, n_{\text{N}_2\text{O}_4}} \quad \text{partial molar Gibbs free energy or chemical potential}$$

\bar{G}_{NO_2} molar Gibbs free energy of **pure** NO_2 $\left[\bar{G}_{\text{NO}_2} \equiv (\Delta \bar{G}_f)_{\text{NO}_2} \right]$

μ_{NO_2} molar Gibbs free energy of NO_2 **in environment where other molecules are present**

thermodynamics of multicomponent systems
E&R section 6.4 (later)

6

$$\mu_i \approx \bar{G}_i$$

for now $\mu_i \approx \bar{G}_i$

$$\Delta\mu_{\text{reaction}} \approx \Delta G_{\text{reaction}}$$

$$\Delta\mu_{\text{reaction}} = \sum_i \nu_i \mu_i \approx \sum_i \nu_i \bar{G}_i = \Delta G_{\text{reaction}}$$

7

7. $\Delta G_{\text{reaction}}$ as a function of pressure, concentration

7. How does $\Delta G_{\text{reaction}}$ ($\Delta\mu$) vary as the concentration of reactants and products varies?

example: 'concentration' of gas = partial pressure P_i

$$P_i = X_i P_{\text{total}} \quad \text{where } X_i \text{ is mole fraction of species } i$$

$$d\bar{G} = -\bar{S}dT + \bar{V}dP \Rightarrow \left(\frac{\partial \bar{G}}{\partial P}\right)_T = \bar{V}$$

for ideal gas:

$$P_{\text{total}} = n_{\text{total}} \frac{RT}{V}$$

$$P_i = n_i \frac{RT}{V}$$

8

7.1 $\Delta G_{\text{reaction}}$ as a function of concentration

$$\left(\frac{\partial \bar{G}}{\partial P}\right)_T = \bar{V}$$

for ideal gas

$$P_{\text{total}} = n_{\text{total}} \frac{RT}{V}$$

$$P_i = n_i \frac{RT}{V}$$

$$\left(\frac{\partial \bar{G}_i}{\partial P_i}\right)_T = \bar{V}$$

$$\bar{G}_i(P_i) - \bar{G}_i(P_i = 1 \text{ bar}) = \int_{1 \text{ bar}}^{P_i} \left(\frac{\partial \bar{G}_i}{\partial P_i'}\right)_T dP_i' = \int_{1 \text{ bar}}^{P_i} \bar{V} dP_i' = \int_{1 \text{ bar}}^{P_i} \frac{RT}{P_i'} dP_i'$$

$$\bar{G}_i(P_i) - \bar{G}_i(P_i = 1 \text{ bar}) = RT \ln\left(\frac{P_i}{1 \text{ bar}}\right)$$

9

7.2 $\Delta G_{\text{reaction}}$ as a function of concentration (ideal gas)

$$\bar{G}_i(P_i) - \bar{G}_i(P_i = 1 \text{ bar}) = RT \ln\left(\frac{P_i}{1 \text{ bar}}\right)$$

$$\bar{G}_i(P_i) - \bar{G}_i^0 = RT \ln\left(\frac{P_i}{1 \text{ bar}}\right)$$

$$\bar{G}_i(P_i) = \bar{G}_i^0 + RT \ln\left(\frac{P_i}{1 \text{ bar}}\right)$$

$$\text{later } \mu_i(P_i) = \mu_i^0 + RT \ln\left(\frac{P_i}{1 \text{ bar}}\right)$$

10

Chemistry 163B Winter 2020
Lecture 15- Free Energy and Equilibrium

7.3 $\Delta G_{\text{reaction}}$ as a function of concentration

$$\bar{G}_i(P_i) = \bar{G}_i^0 + RT \ln\left(\frac{P_i}{1 \text{ bar}}\right)$$

$$\Delta G_{\text{reaction}} = \sum_i \nu_i \bar{G}_i$$

$$\Delta G_{\text{reaction}} = \sum_i \nu_i \left(\bar{G}_i^0 + RT \ln\left(\frac{P_i}{1 \text{ bar}}\right) \right)$$

$$\Delta G_{\text{reaction}} = \sum_i \nu_i \bar{G}_i^0 + \sum_i \nu_i RT \ln\left(\frac{P_i}{1 \text{ bar}}\right)$$

$$\Delta G_{\text{reaction}} = \Delta G_{\text{reaction}}^0 + \sum_i \nu_i RT \ln\left(\frac{P_i}{1 \text{ bar}}\right)$$

$$\underline{RT} \sum_i \bar{\nu}_i \ln\left(\frac{P_i}{1 \text{ bar}}\right) = \underline{RT} \ln \left[\prod_i \left(\frac{P_i}{1 \text{ bar}}\right)^{\bar{\nu}_i} \right] = \underline{RT} \ln Q_P$$

$$Q_P = \prod_i \left(\frac{P_i}{1 \text{ bar}}\right)^{\bar{\nu}_i}$$

for 'persnickety' units notation

$$\underline{R} = R(\text{mol}) = [J K^{-1}]$$

$$\bar{\nu}_i = \nu_i(\text{mol}^{-1}) = [\text{unitless}]$$


11

7.4 $\Delta G_{\text{reaction}}$ as a function of concentration

$$\Delta G_{\text{reaction}} = \Delta G_{\text{reaction}}^0 + \underline{RT} \ln Q_P$$

$$\text{reaction quotient } Q_P = \prod_i \left(\frac{P_i}{1 \text{ bar}}\right)^{\bar{\nu}_i} = \frac{\prod_{q=\text{prod}} \left(\frac{P_q}{1 \text{ bar}}\right)^{\bar{\nu}_q}}{\prod_{r=\text{react}} \left(\frac{P_r}{1 \text{ bar}}\right)^{\bar{\nu}_r}} \quad \mathbf{Q \text{ is UNITLESS}}$$

1 bar is P^0 , std state for P



$$Q_P = \frac{\left(\frac{P_C}{1 \text{ bar}}\right)^{\bar{\nu}_c} \left(\frac{P_D}{1 \text{ bar}}\right)^{\bar{\nu}_d}}{\left(\frac{P_A}{1 \text{ bar}}\right)^{\bar{\nu}_a} \left(\frac{P_B}{1 \text{ bar}}\right)^{\bar{\nu}_b}} \quad \text{'like' an equilibrium constant}$$

$$\text{for solutes in soln } Q_c = \prod_i \left(\frac{[i]}{1 \text{ M}}\right)^{\bar{\nu}_i} \quad Q_c = \frac{\left(\frac{[C]}{1 \text{ M}}\right)^{\bar{\nu}_c} \left(\frac{[D]}{1 \text{ M}}\right)^{\bar{\nu}_d}}{\left(\frac{[A]}{1 \text{ M}}\right)^{\bar{\nu}_a} \left(\frac{[B]}{1 \text{ M}}\right)^{\bar{\nu}_b}} \quad \mathbf{Q \text{ is UNITLESS}}$$

HW6 prob 39 **1 M is std state for solute 12**

Chemistry 163B Winter 2020
Lecture 15- Free Energy and Equilibrium

7.5 Q and units (persnicketyness)

$$\Delta G_{\text{reaction}} = \Delta G_{\text{reaction}}^0 + RT \ln Q_p$$

reaction quotient $Q_p = \prod_i \left(\frac{P_i}{1 \text{ bar}} \right)^{\nu_i}$ **Q is UNITLESS**

$$Q_c = \prod_i \left(\frac{[i]}{1 M} \right)^{\nu_i}$$

$$\Delta G_{\text{reaction}} = \Delta G_{\text{reaction}}^0 + \sum_i \nu_i RT \ln \left(\frac{P_i}{1 \text{ bar}} \right)$$

$$\Delta G_{\text{reaction}} = \Delta G_{\text{reaction}}^0 + \sum_i \frac{\nu_i}{1 \text{ mol}} (1 \text{ mol } R) T \ln \left(\frac{P_i}{1 \text{ bar}} \right)$$

$$\Delta G_{\text{reaction}} = \Delta G_{\text{reaction}}^0 + \sum_i (1 \text{ mol } R) T \ln \left(\frac{P_i}{1 \text{ bar}} \right)^{\frac{\nu_i}{1 \text{ mol}}}$$

$$\Delta G_{\text{reaction}} = \Delta G_{\text{reaction}}^0 + \sum_i (R) T \ln \left(\frac{P_i}{1 \text{ bar}} \right)^{\nu_i} \quad \leftarrow \text{unitless exponent}$$

$$[R \times 1 \text{ mol}] = [R] = J K^{-1}$$

13

7.6 $\Delta G_{\text{reaction}}$ as a function of concentration

$$\Delta G_{\text{reaction}} = \Delta G_{\text{reaction}}^0 + RT \ln Q$$

$$Q_p = \prod_i \left(\frac{P_i}{1 \text{ bar}} \right)^{\nu_i} \quad Q_c = \prod_i \left(\frac{[i]}{1 M} \right)^{\nu_i}$$

evaluates ΔG for **ANY** set of pressures, concentrations

ΔG^0 gives free energy change for standard conditions,
 $RT \ln Q$ corrects ΔG for actual P's and [conc's]

at equilibrium $\Delta G = ?$

14

8. $\Delta G_{\text{reaction}}$ at equilibrium

at equilibrium $\Delta G = 0$

$$0 = \Delta G_{\text{reaction}}^0 + \underline{RT} \ln Q_{\text{eq}}$$

$$\Delta G_{\text{reaction}}^0 = -\underline{RT} \ln Q_{\text{eq}}$$

at given T $Q_{\text{eq}} = \text{constant} \equiv K_{\text{eq}}$

$$\Delta G^0 = -\underline{RT} \ln K_{\text{eq}}$$

$$K_{\text{eq}} = e^{-\frac{\Delta G^0}{RT}}$$

$$\Delta G_{\text{reaction}} = \Delta G_{\text{reaction}}^0 + \underline{RT} \ln Q$$

$$Q_p = \prod_i \left(\frac{P_i}{1 \text{ bar}} \right)^{\nu_i} \quad Q_c = \prod_i \left(\frac{[i]}{1 M} \right)^{\nu_i}$$


$\Delta G_{\text{reaction}} = \Delta G_{\text{reaction}}^0 + \underline{RT} \ln Q$ any concentrations for Q

$Q_{\text{eq}} \equiv K_{\text{eq}}$ for equilibrium concentrations

that satisfy $\Delta G_{\text{reaction}}^0 = -\underline{RT} \ln K_{\text{eq}}$

15

9. $\Delta G_{\text{reaction}} = \Delta G^0 + \underline{RT} \ln Q$ is extensive



?? $\Delta G_2 = 2\Delta G_1$??

$$Q_1 = \frac{\left(\frac{P_C}{1 \text{ bar}} \right)^{\bar{n}_c} \left(\frac{P_D}{1 \text{ bar}} \right)^{\bar{n}_d}}{\left(\frac{P_A}{1 \text{ bar}} \right)^{\bar{n}_a} \left(\frac{P_B}{1 \text{ bar}} \right)^{\bar{n}_b}} \quad Q_2 = \frac{\left(\frac{P_C}{1 \text{ bar}} \right)^{2\bar{n}_c} \left(\frac{P_D}{1 \text{ bar}} \right)^{2\bar{n}_d}}{\left(\frac{P_A}{1 \text{ bar}} \right)^{2\bar{n}_a} \left(\frac{P_B}{1 \text{ bar}} \right)^{2\bar{n}_b}}$$

$$Q_2 = Q_1^2$$

$$\Delta G_1 = \Delta G_1^0 + \underline{RT} \ln Q_1$$

$$\Delta G_2 = \Delta G_2^0 + \underline{RT} \ln Q_2$$

$$\Delta G_2 = 2\Delta G_1^0 + \underline{RT} \ln Q_1^2 = 2(\Delta G_1^0 + \underline{RT} \ln Q_1)$$

$$\Delta G_2 = 2\Delta G_1$$

16

10.1 variation of K_{eq} with T

- $(\Delta G_{rxn})_{T,P} \Rightarrow$ reaction carried out isothermally a P
- Vary T: $(\Delta G_{rxn})_{T_1,P}$ vs $(\Delta G_{rxn})_{T_2,P}$
- Need $\left(\frac{\partial G}{\partial T}\right)_P$ and $\left(\frac{\partial \Delta G_{rxn}}{\partial T}\right)_P$

17

10.2 variation of K_{eq} with T

$$d\bar{G} = -\bar{S}dT + \bar{V}dP$$

$$\left(\frac{\partial \bar{G}}{\partial T}\right)_P = -\bar{S} \quad \text{but remember } S(T)$$

a few manipulations which lead to simpler final relationships

$$\Delta G_{rxn} = \Delta G_{rxn}^0 + RT \ln Q \quad \Delta G_{rxn}^0 = -RT \ln K_{eq}$$

$$\frac{\Delta G}{T} = \frac{\Delta G^0}{T} + R \ln Q \quad \frac{\Delta G^0}{T} = -R \ln K_{eq}$$

$$\left(\frac{\partial \frac{\Delta G}{T}}{\partial T}\right)_P = \frac{1}{T} \left(\frac{\partial \Delta G}{\partial T}\right)_P - \frac{\Delta G}{T^2}$$

18

10.3 variation of K_{eq} with T

$$\left(\frac{\partial \frac{\Delta G}{T}}{\partial T}\right)_P = \frac{1}{T} \left(\frac{\partial \Delta G}{\partial T}\right)_P - \frac{\Delta G}{T^2}$$

$$\left(\frac{\partial \frac{\Delta G}{T}}{\partial T}\right)_P = -\frac{\Delta S}{T} - \frac{\Delta G}{T^2}$$

$$\left(\frac{\partial \frac{\Delta G}{T}}{\partial T}\right)_P = -\frac{\Delta S}{T} - \frac{(\Delta H - T\Delta S)}{T^2}$$

$$\boxed{\left(\frac{\partial \frac{\Delta G}{T}}{\partial T}\right)_P = -\frac{\Delta H}{T^2}}$$

19

10.4 variation of K_{eq} with T

$$\left(\frac{\partial \frac{\Delta G}{T}}{\partial T}\right)_P = -\frac{\Delta H}{T^2}$$

$$\left(\frac{\partial \frac{\Delta G^0}{T}}{\partial T}\right)_P = \left(\frac{\partial (-R \ln K)}{\partial T}\right)_P = -R \left(\frac{\partial \ln K}{\partial T}\right)_P = -\frac{\Delta H^0}{T^2}$$

$$\boxed{\left(\frac{\partial \ln K}{\partial T}\right)_P = +\frac{\Delta H^0}{RT^2}}$$

20

10.5 variation of K_{eq} with T

$$\left(\frac{\partial \ln K}{\partial T}\right)_p = +\frac{\Delta H^0}{RT^2}$$

$$\int_{T_1}^{T_2} d \ln K = \int_{T_1}^{T_2} +\frac{\Delta H^0}{RT^2} dT$$

$$\ln K_{T_2} - \ln K_{T_1} = -\frac{\Delta H^0}{R} \left(\frac{1}{T_2} - \frac{1}{T_1}\right) = \frac{\Delta H^0}{R} \left(\frac{1}{T_1} - \frac{1}{T_2}\right)$$

$$\ln \left(\frac{K_{T_2}}{K_{T_1}}\right) = \frac{\Delta H^0}{R} \left(\frac{1}{T_1} - \frac{1}{T_2}\right)$$

21

10.6 variation of K_{eq} with T

$$\ln \left(\frac{K_{T_2}}{K_{T_1}}\right) = \frac{\Delta H^0}{R} \left(\frac{1}{T_1} - \frac{1}{T_2}\right)$$

$$T_2 > T_1$$


endothermic, $\Delta H^0 > 0 \Rightarrow \ln \frac{K_{T_2}}{K_{T_1}} > 0 \Rightarrow K_{T_2} > K_{T_1}$

higher T moves equilibrium to right (products) reactants + heat \rightleftharpoons products

exothermic $\Delta H^0 < 0 \Rightarrow \ln \frac{K_{T_2}}{K_{T_1}} < 0 \Rightarrow K_{T_2} < K_{T_1}$

higher T moves equilibrium to left (reactants) reactants \rightleftharpoons products + heat

Le Chatelier's Principle
"for heat"



22

goals of lecture

- ✓ 6. Brief hello to thermodynamics of multicomponent systems (n's vary)
- ✓ 7. $\Delta G_{\text{reaction}}$ for non-standard state concentrations, pressures
$$\Delta G_{\text{reaction}} = \Delta G^{\circ}_{\text{reaction}} + RT \ln Q$$
- ✓ 8. K_{eq} and $\Delta G^{\circ}_{\text{reaction}}$
- ✓ 9. $\Delta G_{\text{reaction}} = \Delta G^{\circ}_{\text{reaction}} + RT \ln Q$ is extensive
- ✓ 10. Variation of K_{eq} with T

23



free

End of Lecture 15

energy



24

Chemistry 163B Winter 2020
Lecture 15- Free Energy and Equilibrium

persnickity



per·snick·et·y

/pərˈsnɪkədē/



Learn to pronounce

adjective INFORMAL • NORTH AMERICAN

placing too much emphasis on trivial or minor details; fussy.
"she's very persnickety about her food"

Similar: fussy difficult to please difficult finicky overfastidious ▾

• requiring a particularly precise or careful approach.



25