

Chemistry 1B

Fall 2016

~~lectures~~ topics 1-2

[ch 12 pp 522-537]_{7th}

- The “laws of nature” in 1900
(successful for describing large objects)
describe particles **AND** describe waves
- Experiments that contradicted these laws
(when applied on the scale of atomic dimensions)
 - ✓ Ultraviolet Catastrophe and Photoelectric Effect
 - ✓ Spectrum of Hydrogen Atom
 - ✓ Davisson-Germer and Compton Experiments
- particles **BEHAVE AS** waves
waves **BEHAVE AS** particles
- obtain and use observed quantitative relationships (HW #1)
- **Why?** To understand the behavior of electrons
in atoms and molecules

- fundamental **particles** and charge
 - electron: - charge, $m_e = 9.109 \times 10^{-31}$ kg
 - proton: + charge, $m_p = 1.672 \times 10^{-27}$ kg
 - neutron: 0 charge, $m_n = 1.675 \times 10^{-27}$ kg


[back page]

- **particles** in general



- electromagnetic **waves**

light waves





- electromagnetic wave
 - fig 12.1
 - ~ fig 12.2
 - fig 12.3
 - spectrum of visible light
- **wave** phenomena
 - refraction
 - dispersion
 - diffraction
 - interference and Fig. 12.7

- [Blackbody radiation- Fig 12.4](#)
- [Ultraviolet catastrophe](#) (p. 525)

• **$E = h\nu$ (energy per photon)**

$h = \text{constant} = 6.626 \times 10^{-34} \text{ m}^2 \text{ kg/s}$ [J-sec]
[J]=[E]=[kg m² s⁻²]

(HW#1 PROB 12.21)



- kinetic energy
- potential energy
- conservation of energy
- momentum (p. 158)
- units and conversions

Chemistry IB -AL

Experiments that ushered in the QUANTUM AGE

the Photoelectric Effect

Windows Media Player
Video01_Photoelectric





~~clicker material for AL class lecture session 2~~

Chem 1B-AL Study Guides TWO GOOD REASONS

reason 1: they are designed to encourage you to **ACTIVELY** interact with the material presented

reason 2: the clicker questions at the beginning of class will test your understanding of the material covered in the study guide

Next class I and II!



2

01:23 [media controls]

Learning Objectives and Worksheet I


Chemistry 1B-AL Fall 2016

Lectures (1-2) Nature of Light and Matter, Quantization of Energy, and the Wave-Particle Duality

I. Understand the following aspects of the laws of physics circa 1900, which were successful for describing large objects (i.e. particles) and electromagnetic waves:

1. Be able to identify the following fundamental particles, their respective charges, and their relative masses (*high school review?? learned on MOM's knee in your childhood??*)
 2. What are three properties that were considered by 19th century physicists to be exclusively 'properties of particles'?
 3. What are three characteristics or properties that were considered by 19th century physicists to be exclusively 'properties of waves'?
 4. Waves (sound, water, electromagnetic, etc.) are characterized by several properties
 5. Know the types of radiation found at various frequencies of electromagnetic radiation and their relative ordering (as a function of wavelength)
 6. Visible electromagnetic radiation fall correspond to wavelengths
 7. Understand the following terms, relationships, and phenomena. Which of these were associated with the classical concept of particles, which with classical electromagnetic waves, and which with both (in classical models)? [*most important are in bold*]
 - i. **Kinetic energy**
 - ii. **Potential energy** (e.g. electrostatic pull, gravitation, stretched spring)
 - iii. **Conservation of energy**
 - iv. **Relation between frequency and wavelength**
 - v. **Momentum**
 - vi. Maxwell's Laws (*will study in physics 6; describes electromagnetic waves*)
 - vii. Dispersion
 - viii. Refraction
 - ix. Diffraction
 - x. **Interference** (constructive and destructive)
-

Introduction: Between 1887 and 1927 experiments indicated that the distinctions between particles and waves were less clear cut than previously postulated



II. Understand the following observations and their implications regarding the deficiencies of the classical physics model

1. Photons:

i. Experiment: BLACKBODY RADIATION

a. **What was observed?**



b. **What was the contradiction to laws of classical physics?**

c. **What was the postulate that resolved contradiction?**


Electromagnetic energy was radiated in packets (photons) with fixed energies

related to their wavelength Planck's law $E_{\text{photon}} = \underline{h\nu} = hc/\lambda$.

where $h = 6.626 \times 10^{-34} \text{ m}^2 \text{ kg s}^{-1}$ (or $h = 6.626 \times 10^{-34} \text{ J sec}$) is Planck's constant.

Know how to find/calculate all variables in Planck's equation

Photoelectric effect



Chemistry 1B-AL Fall 2016, Study Guide and Worksheet I

Chemistry 1B-AL Fall 2016, Study Guide and Worksheet I

- c. What was the postulate that resolved contradiction?
 Electromagnetic energy was radiated in packets (photons) with fixed energies related to their wavelength. Planck's law $E_{\text{photon}} = \frac{hc}{\lambda}$, where $h = 6.626 \times 10^{-34} \text{ m}^2 \text{ kg s}^{-1}$ Planck's constant.

d. Know how to find/calculate all variables in Planck's equation

- ii. Experiment: PHOTOELECTRIC EFFECT (Einstein won [Nobel Prize 1921](#) for interpretation of the Photoelectric Effect)

a. What was observed?

b. What was the contradiction to laws of classical physics?

c. What classical physics would predict??

(1) Long wavelength, small amplitude electromagnetic wave

(2) Long wavelength, large amplitude electromagnetic wave

d. What was observed in terms of ejected electrons:

Wavelength of light	Intensity of light	Relative number of electrons (none/few/many)	Relative velocity of electrons (slow/fast)
long ($hc/\lambda < \Phi$)	weak	_____	_____
long ($hc/\lambda < \Phi$)	strong	_____	_____
medium (hc/λ just above Φ)	weak	_____	_____
short ($hc/\lambda \gg \Phi$)	weak	_____	_____
short ($hc/\lambda \gg \Phi$)	strong	_____	_____

e. What was the postulate that resolved contradiction?





HW #1: 3, 4, 5

f. Be able to calculate the minimum energy to remove an electron from a "potential well" and the K.E. of any ejected electrons.

c

photoelectric effect (pp 514-515)

- [apparatus Sil Fig. 7.7](#) →
- [what's observed](#)
- interpretation



"electron in potential well"

Windows Media Player
Video01_Photoelectric_effect

"electron in potential well"

C_s
 $3.05 \times 10^{-19} \text{ J}$

Now
 $3.94 \times 10^{-19} \text{ J}$

Φ is work function

$\Phi_K = 3.59 \times 10^{-19} \text{ J}$

07:34

7

what was expected classically

Windows Media Player

Video01_Photoelectric_effect

what was expected classically

red light

weak



strong, intense



Chemistry 1B-AL Fall 2016, Study Guide and Worksheet 1

c. What was the postulate that resolved contradiction?
Electromagnetic energy was radiated in packets (photons) with fixed energies related to their wavelength. Planck's law $E_{\text{photon}} = \frac{hc}{\lambda}$, where $h = 6.626 \times 10^{-34} \text{ m}^2 \text{ kg s}^{-1}$ Planck's constant.

d. Know how to find/calculate all variables in Planck's equation

ii. Experiment: PHOTOELECTRIC EFFECT (Einstein won Nobel Prize 1921 for interpretation of the Photoelectric Effect)

a. What was observed?

b. What was the contradiction to laws of classical physics?

c. What classical physics would predict??


(1) Long wavelength, small amplitude electromagnetic wave

(2) Long wavelength, large amplitude electromagnetic wave

08:40



photoelectric effect : electron in a “potential well”
(**CLASSICAL**: *small amplitude, long wavelength*)




no electron
ejected

Φ is work function (value depends on the metal being irradiated)

example Φ for potassium= $3.68 \times 10^{-19} \text{J} = 2.29 \text{eV}$

photoelectric effect : electron in a "potential well"
(CLASSICAL: large amplitude, long wavelength)



nada: no electron ejected (RED IS JUST A LOSER WAVELENGTH)
CLASSICALLY:


just increase amplitude to get enough energy to eject electron !!


what was observed


Windows Media Player
Video01_Photoelectric_effect

what was observed

vary λ of incident light

 long λ
low ν

 med λ
med ν


 short λ
high ν

- ① are e^- 's emitted
- ② Do e^- 's have K.E.
- ③ Number of e^- 's vs intensity of light


11

11:25

photoelectric effect : electron in a "potential well" ($\lambda = \text{long}$)



photoelectric effect : electron in a "potential well" ($\lambda = \text{medium}$)




example: for potassium (depth of 'potential well')

$$\Phi = 3.59 \times 10^{-19} \text{ J} = 2.24 \text{ eV}$$


photon of this energy: $E = h\nu = hc/\lambda$ $\lambda = (c/\nu) = (ch/E) = 553 \text{ nm}$

($\lambda = 553 \text{ nm}$ color?)

photoelectric effect : electron in a "potential well" ($\lambda = \text{short}$)



summary of observations



for given $\nu > \nu_0$ increase intensity of light

⇒ more photons, but E per photon remains same

⇒ more electrons, but of same kinetic energy

conservation of energy (p 527)

if an individual photon does not have sufficient energy to 'kick' electron out of the potential well of metal:

NO ELECTRONS EJECTED !!

DEFINITION work function of metal : $\Phi_{\text{metal}} \equiv h\nu_0 = hc/\lambda_0$ i.e. depth of well and the minimum energy to eject electron

if a individual photon that has sufficient energy to 'kick' electron out of the potential well of metal:

energy of photon = energy to get out of well + kinetic energy of electron


$$h\nu_{\text{photon}} = h\nu_0 + \text{K.E.} = \Phi_{\text{metal}} + \frac{1}{2} m_e v_{\text{electron}}^2$$

$$\text{K.E.} = h\nu_{\text{photon}} - h\nu_0 = h\nu_{\text{photon}} - \Phi_{\text{metal}}$$

$$v_{\text{electron}} = \left(\frac{2}{m_e} (h\nu_{\text{photon}} - \Phi_{\text{metal}}) \right)^{1/2}$$

(HW#1 12.27,
12.29)

class activity on photoelectron emission



Next Class Session !!!

2. Energies of electrons in atoms are quantized:
 3. “Particles” behaving like “waves”
 4. “Waves” behaving like “particles” (actually “more of waves behaving like particles” since the quanta of radiation in the photoelectric effect may be thought to be examples of light with particle-like behavior)
- III. Measurement and the Uncertainty Principle (consequences of wave-like properties)

Finis For Today

- classical prediction ([death spiral](#))




- observation of atomic spectra
[fig. 12.8](#)

$$\frac{1}{\lambda} = R \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \quad n_2 \text{ and } n_1 \text{ are integers with } n_2 > n_1$$

$R = 1.096776 \times 10^7 \text{ m}^{-1}$ R is the Rydberg constant

- Bohr model (HW#1 12.38a,d, 12.40)
[fig. 12.10](#), [Silberberg 7.10](#)






Dr. Quantum Pair-Share Activity



Davisson-Germer experiment

remember: WAVES showed constructive and destructive interference





diffraction by slits and crystals




mind blowing

X-ray diffraction
destructive int
of Al foil

destructive int
Electron diffraction
of Al foil

- **diffraction of electrons-**
(Davisson-Germer Experiment; p. 530)
- **De Broglie relationship (p. 528)** (HW#1 12.32,12.35)
- **What is “meaning” of electron wave??**
(<http://phys.educ.ksu.edu/vqm/html/doubleslit/index.html>) 
(<http://www.youtube.com/watch?v=DfPeprQ7oGc>) 
- **Wavelengths of “ordinary” objects**
(p. 528, example 12.2) Silberberg Table 7.1 (HW#1 prob S2) 
- **Compton Experiment**
- Heisenberg uncertainty principle (p. 539) *boing!!* (HW#1 12.49) 

- ✓ • The “laws of nature” in 1900
(successful for describing large objects)
describe particles **AND** describe waves
- ✓ • Experiments that contradicted these laws
(when applied on the scale of atomic dimensions)
 - ✓ ✓ Davisson-Germer and Compton Experiments
 - ✓ ✓ Ultraviolet Catastrophe and Photoelectric Effect 
 - ✓ ✓ Spectrum of Hydrogen Atom
- ✓ • particles **BEHAVE AS** waves
waves **BEHAVE AS** particles
- ✓ • Ensuing quantitative relationships

What to do ??

invent quantum mechanics !!!

Solvay Conference 1927




The mid-1920's saw the development of the quantum theory, which had a profound effect on chemistry. Many theories in science are first presented at international meetings. This photograph of well-known scientists was taken at the international Solvay Conference in 1927. Among those present are many whose names are still known today. Front row, left to right: I. Langmuir, **M. Planck**, **M. Curie**, H. A. Lorentz, **A. Einstein**, P. Langevin, C. E. Guye, C. T. R. Wilson, O. W. Richardson. Second row, left to right: P. Debye, M. Knudsen, W. L. Bragg, H. A. Kramers, P. A. M. Dirac, A. H. Compton, **L. V. de Broglie**, M. Born, **N. Bohr**. Standing, left to right: A. Piccard, E. Henriot, P. Ehrenfest, E. Herzen, T. De Donder, **E. Schroedinger**, E. Verschaffelt, **W. Pauli**, **W. Heisenberg**, R. H. Fowler, L. Brillouin.

'fuel' for quantum mechanics




Werner Heisenberg and Niels Bohr dining in Copenhagen in 1934.



end of topics1-2

figure 12.1 Zumdahl




©Houghton Mifflin Company. All rights reserved.




wavelength (λ) and frequency (ν)

wave amplitude

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



HW PROBS #12.22, 12.23




$$\lambda \nu = c$$


c = speed of light
 $2.996 \times 10^8 \text{ m s}^{-1}$

Silberberg figure 7.2

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Zumdahl figure 12.3




©Houghton Mifflin Company. All rights reserved.

HW PROBS #12.21, #12.23



wavelength and color





R O Y G B I V

ROY G BIV



Silberberg figure 7.5 and Zumdahl 12.7


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




©Houghton Mifflin Company. All rights reserved.



Zumdahl figure 12.3 and Silberberg figure 7.6



©Houghton Mifflin Company. All rights reserved.



Smoldering coal



Electric heating element




Lightbulb filament




The Ultraviolet Catastrophe

Unfortunately, the theory disagree violently with experiment





Silberberg figure 7.7

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Zumdahl figure 12.8



(b)



BUT ???




Figure 17–9 In classical theory, atoms constructed according to the Rutherford model are not stable; the electron would quickly spiral into the nucleus.



Silberberg figure 7.10 (emission: H-electron loses energy)

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

$$E_n = -2.178 \times 10^{-18} \text{ J} \left(\frac{Z^2}{n^2} \right) \quad Z=1 \text{ for H atom, } n=1, 2, 3, \dots$$

energy lost (emitted) by H atom, $n_2 \rightarrow n_1$ ($n_2 > n_1$)

$$\Delta E_{n_2 \rightarrow n_1} = E_{n_1} - E_{n_2} = -2.178 \times 10^{-18} \text{ J } Z^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$


energy of photon emitted

$$E_{\text{photon}} = -\Delta E_{n_2 \rightarrow n_1}$$

$$h\nu_{\text{photon}} = 2.178 \times 10^{-18} \text{ J } Z^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

$$\frac{hc}{\lambda_{\text{photon}}} = 2.178 \times 10^{-18} \text{ J } Z^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

$$\frac{1}{\lambda} = \frac{2.178 \times 10^{-18} \text{ J}}{hc} Z^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) = 1.0967 \times 10^7 \text{ m}^{-1} Z^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$



Silberberg figure 7.10 (absorption, H-electron gains energy)

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

$$E_n = -2.178 \times 10^{-18} \text{ J} \left(\frac{Z^2}{n^2} \right) \quad Z=1 \text{ for H atom, } n=1, 2, 3, \dots$$

energy gained (absorbed) by H atom, $n_1 \rightarrow n_2$ ($n_2 > n_1$)

$$\Delta E_{n_1 \rightarrow n_2} = E_{n_2} - E_{n_1} = -2.178 \times 10^{-18} \text{ J } Z^2 \left(\frac{1}{n_2^2} - \frac{1}{n_1^2} \right)$$


energy of photon absorbed

$$E_{\text{photon}} = +\Delta E_{n_1 \rightarrow n_2}$$


$$h\nu_{\text{photon}} = 2.178 \times 10^{-18} \text{ J } Z^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \text{ (note loss of - and switch of } n_1, n_2 \text{)}$$

$$\frac{hc}{\lambda_{\text{photon}}} = 2.178 \times 10^{-18} \text{ J } Z^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$


$$\frac{1}{\lambda} = \frac{2.178 \times 10^{-18} \text{ J}}{hc} Z^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) = 1.0967 \times 10^7 \text{ m}^{-1} Z^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$



Zumdahl figure 12.10



(a)




(c)

©Houghton Mifflin Company. All rights reserved.




Silberberg figure 7.5 and Zumdahl 12.7


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




(a) Diffraction



(b) Constructive interference



(c) Destructive interference



$$\lambda = \frac{h}{m_e v_e}$$



photons $\lambda = 402 \text{ nm}$

λ increases $\approx \times 1.5$



photons $\lambda = 594 \text{ nm}$

[slit 10^{-5} m]



electrons $v = 4.6 \times 10^6 \text{ m sec}^{-1}$

v decreases $\approx \times 1.5$



electrons $v = 3.06 \times 10^6 \text{ m sec}^{-1}$

[slit $5 \times 10^{-9} \text{ m}$]



Silberberg Table 7.1

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

**note mass in g, need to use kg for $mv\lambda=h$
(λ correct in table)**

Substance	Mass (g)	Speed (m/s)	λ (m)
Slow electron	9×10^{-28}	1.0	7×10^{-4}
Fast electron	9×10^{-28}	5.9×10^6	1×10^{-10}
Alpha particle	6.6×10^{-24}	1.5×10^7	7×10^{-15}
One-gram mass	1.0	0.01	7×10^{-29}
Baseball	142	25.0	2×10^{-34}
Earth	6.0×10^{27}	3.0×10^4	4×10^{-63}



letters to nature

NATURE | VOL 401 | 14 OCTOBER 1999 |

.....

Wave-particle duality of C_{60} molecules


**Markus Arndt, Olaf Nairz, Julian Vos-Andreae, Claudia Keller,
Gerbrand van der Zouw & Anton Zeilinger**

*Institut für Experimentalphysik, Universität Wien, Boltzmannngasse 5,
A-1090 Wien, Austria*

.....



waves: Δx vs Δv




Δx

Interactive Java Tutorials

Diffraction

Modified version of

<http://www.micro.magnet.fsu.edu/primer/java/scienceopticsu/diffraction/basicdiffraction/index>



Δx larger


Δv_x smaller

Interactive Java Tutorials

Diffraction

Modified version of

<http://www.micro.magnet.fsu.edu/primer/java/scienceopticsu/diffraction/basicdiffraction/index>



Δx smaller

Δv_x larger



Uncertainty and measurement





FIGURE 15.24 A photon, which has a negligible effect on the trajectory of a baseball (a), significantly perturbs the trajectory of the far less massive electron (b).



- particles have mass (m), definite positions (x) and velocities (v)



- particles have kinetic energy and momentum $KE = \frac{1}{2}mv^2$
 $p = mv$



- particles obey Newton's laws of physics $F = ma$



CHEMISTRY 1B-AL FALL 2016

HANDOUTS

COURSE INFORMATION	TOPICS SCHEDULE	HOMEWORK ASSIGNMENTS	LEARNING OBJECTIVES WORKSHEETS & VIDEOS	DISCUSSION SECTIONS
GALLERY OF MOLECULES JMOL	HANDOUTS	TEAM BASED ACTIVE LEARNING	HELP SCHEDULE	TO ECOMMONS
				ON-LINE Q & A

go to current handouts

	HANDOUT	DATE NEEDED	FORMAT
1	WebAssign LogIn Access	23 Sept	PDF
2	WebAssign LogIn Test	23 Sept	PDF
3	iClicker Registration	23 Sept	PDF
4	Homework #1	23 Sept	PDF
5	Topic Handouts (1-2)	23 Sept	PDF(6) PDF(2) PDF(1)



7	Summary of Experiments leading to Quantum Theory	28 Sept	PDF
---	--	---------	-----

REVIEW

The enigma of optical momentum in a medium

BY STEPHEN M. BARNETT^{1,*} AND RODNEY LOUDON²

¹*Department of Physics, SUPA, University of Strathclyde,
Glasgow G4 0NG, UK*

²*School of Computer Science and Electronic Engineering, University of Essex,
Colchester CO4 3SQ, UK*

It is 100 years since Minkowski and Abraham first gave rival expressions for the momentum of light in a material medium. At the single-photon level, these correspond, respectively, either to multiplying or dividing the free-space value ($\hbar\mathbf{k}$) by the refractive index (n). The debate that this work started has continued till the present day, punctuated by the occasional publication of ‘decisive’ experimental demonstrations supporting one or other of these values. We review the compelling arguments made in support of the Minkowski and Abraham forms and are led to the conclusion that *both* momenta are correct. We explain why two distinct momenta are needed to describe light in a medium and why each appears as the natural, and experimentally observed, momentum in appropriate situations.

Keywords: Abraham–Minkowski dilemma; photon momentum; Poynting vector;
quantum optics
