

# Crown 85: Visual Perception: A Window to Brain and Behavior



Lecture 3: Techniques for Studying Brain and Behavior

# lecture 3 outline

---

Crown 85 Winter 2016

Visual Perception: A Window to Brain and Behavior

Lecture 3- Techniques for Studying the Brain

Reading: [Amherst College Course](#)  
[Stanford College Course](#)

Looking: [Optogenetics](#)  
[ECoG](#)

Understand the basic principles upon which the following techniques and the kinds of information about neural processing which their application can provide:

*disclaimer !!!!!*

---

Just a



not expecting U to become an



© Can Stock Photo

# 1. Anatomical

a. Neuron staining

b. Electron microscopy

c. Pathway tracing


# Anatomy: Golgi, Nissl staining of neural substrate

---




Camillo Golgi  
1843-1926

**Golgi**  
staining of **entire neuron** with  
silver chromate.  
only stains a subset of cells



<http://www.nature.com/npp/journal/v32/n10/covers/index.html>

# Anatomy: Golgi, Nissl staining of neural substrate



Franz Nissl  
1860-1919

**Nissl**  
staining of **cell body** with dyes  
(e.g. cresyl violet) that interact  
with RNA (and DNA) .




Figure 13. Nissl stain of the visual cortex reveals the different layers I through VI quite clearly.

<http://webvision.med.utah.edu/imageswv/nissl2.jpg>



The Nobel Prize in Physiology or Medicine 1906

Camillo Golgi, Santiago Ramón y Cajal

Share this:

## The Nobel Prize in Physiology or Medicine 1906



Camillo Golgi  
Prize share: 1/2



Santiago Ramón y Cajal  
Prize share: 1/2

The Nobel Prize in Physiology or Medicine 1906 was awarded jointly to Camillo Golgi and Santiago Ramón y Cajal *"in recognition of their work on the structure of the nervous system"*

Photos: Copyright © The Nobel Foundation


OPINION

## The contribution of Santiago Ramón y Cajal to functional neuroscience

Rodolfo R. Llinás

Santiago Ramón y Cajal — arguably the most accomplished anatomist in the history of neuroscience — became recognized as such not only because of his incredible anatomical skills and his indefatigable working habits, but also because of his uncanny sense of the functional implications of his work, a sense that made him a true genius in the field of biology

<http://www.utdallas.edu/~tres/memory/intro/llinas.pdf>



**Figure 33.19**  
*Biochemistry, Seventh Edition*  
© 2012 W. H. Freeman and Company

## **Figure 32.20** The Rod Cell

(Left) Scanning electron micrograph of retinal rod cells. (Right) Schematic representation of a rod cell. [Photograph courtesy of Dr. Deric Bownds.]

<http://www.ncbi.nlm.nih.gov/books/NBK22541/>





# *anatomy: tract tracing (e.g. HRP, <sup>3</sup>H proline, etc)*

## Pathway Tracing:

Injecting, via fine needle, a 'tracer substance' into or near a neuron which is then transported down the axon (anterograde: *soma* → *axon terminal*) or up the axon (retrograde : *axon terminal* → *soma*). The pathway is then visualized by the color or radiographic 'footprint' of the tracer. The color may come from a tracer that is itself a dye or one that is produced by a subsequent 'developing' reaction.





A Schematic diagram of dual transsynaptic tracing



<http://journal.frontiersin.org/article/10.3389/neuro.01.032.2009/full>

# *anatomy: tract tracing (e.g. HRP, $^3\text{H}$ proline, etc)*

---



<http://vision.ucsf.edu/hortonlab/ResearchProgram.html>

<http://www.nature.com/nrn/journal/v9/n6/full/nrn2391.html>

<https://www.hccfl.edu/media/153423/handbook%20of%20chemical%20neuroanatomy%20vol8%20ch5.pdf>



1. Anatomical

- ✓ a. Neuron staining
- ✓ b. Electron microscopy
- ✓ c. Pathway tracing

2. Electrophysiological recording of neural activity
  - a. Single cell recording in neurons
  - b. Electroencephalography (EEG)
  - c. Magnetoencephalography (MEG)
  - d. Electrocochicography (ECoG)

# electrophysiology: single cell recordings of neuronal activity



Place an electrode on a single neuron and measure the frequency of firing.



<http://www.kyb.tuebingen.mpg.de/nc/de/mitarbeiter/details/theofanis.html>

## Single-unit recording


stimulus presented on TV screen



visual cortex

lateral geniculate nucleus

recording electrode



direction of motion

neuron's response

# UCSC multielectrode array



ELSEVIER


Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Nuclear Instruments and Methods in Physics Research A 501 (2003) 298–307

**NUCLEAR  
INSTRUMENTS  
& METHODS  
IN PHYSICS  
RESEARCH**  
Section A

[www.elsevier.com/locate/nima](http://www.elsevier.com/locate/nima)



## Large-scale imaging of retinal output activity ☆

A.M. Litke<sup>a,\*</sup>, E.J. Chichilnisky<sup>b</sup>, W. Dabrowski<sup>c</sup>, A.A. Grillo<sup>a</sup>, P. Grybos<sup>c</sup>,  
S. Kachionine<sup>a</sup>, M. Rahman<sup>d</sup>, G. Taylor<sup>a</sup>



<sup>a</sup> Santa Cruz Institute for Particle Physics, University of California, Santa Cruz, CA 95064, USA

<sup>b</sup> The Salk Institute, La Jolla, CA 92037, USA

<sup>c</sup> Faculty of Physics and Nuclear Techniques, University of Mining and Metallurgy, Krakow, Poland

<sup>d</sup> University of Glasgow, Glasgow, UK




 **Crown 85- Visual Perception:**  
A Window to Brain and Behavior 

HOME COURSE VOICER RESEARCH LAB  
INFORMATION LECTURES REPORTS INTERVIEWS PROJECTS eCOMMONS



One goal of Crown 85 is to introduce its students to researchers and research in the various neuroscience and perception laboratories throughout the UCSC campus. Enrolled students are required to visit one of these researchers and present a class report on the studies being conducted in their laboratory. Laboratories which have agreed to host Crown 85 student interviews include:

Prof. David Feldheim Department of Molecular and Cellular Biology	Development of the Visual System	<a href="#">read more</a>
Prof. Alexander 'Sasha' Sher Department of Physics and SCIPP	Retinal Structure and Function Retinal Regeneration Advanced Multi-electrode Recording Techniques	<a href="#">read more</a>



# electrophysiology (neural recordings): electrocorticography (ECoG)

Subdural electrodes implanted inside the cranium. For purely clinical reasons, patients with epilepsy are sometimes implanted with such electrodes to localize their seizure onset prior to surgical therapy.



[https://en.wikiversity.org/wiki/Blausen\\_gallery\\_2014](https://en.wikiversity.org/wiki/Blausen_gallery_2014)


<https://www.med.nyu.edu/thesenlab/research-0/intracranial-eeG/>



# *electrophysiology (neural recordings): electrocorticography (ECoG)*

---

## STANFORD TEAM HELPS WOMAN BATTLE EPILEPTIC SEIZURES





Surgeons at Stanford Hospital helped a Bay Area woman overcome intractable epilepsy and have fewer seizures.

<http://abc7news.com/archive/9405444/>

# electrophysiological recording: electroencephalography (EEG)

**EEG:** measures electric fields produced by neural electrical activity



The time course of shape discrimination in the human brain  
Ales JM, Appelbaum LG, Cottereau BR, Norcia AM

<http://www.st-andrews.ac.uk/~jma23/papers/alesInPressNeuroimageShape.pdf>

<http://www.ski.org/lab/hou-lab>

# *electrophysiological recording: magnetoencephalography (MEG)*

---

**MEG:** measures the magnetic fields generated by neural activity




Image 2 of 8

**SQUID SENSOR ARRAY**



<http://ilabs.washington.edu/what-magnetoencephalography-meg>

<http://www.washington.edu/news/author/mollywmc/>

2. Electrophysiological recording of neural activity
  - ✓ a. Single cell recording in neurons
  - ✓ b. Electroencephalography (EEG)
  - ✓ c. Magnetoencephalography (MEG)
  - ✓ d. Electroconvulsive therapy (ECT)

### 3. Imaging

a. Positron emission tomography (PET)

b. Functional magnetic resonance imaging (fMRI)

c. Calcium dyes

d. Voltage sensitive dyes

# imaging: tomography (CT, PET, OCT)

collection of absorbed or emitted radiation (or electrons/positrons) from multiple detectors distributed in space; allows one to localize and image the source of the absorption or emission





Figure 7-10 Computer tomography

<http://130.237.83.53/medicaldevices/album/Ch%207%20Medical%20images/slides/F%207-10%20Computer%20tomography.html>



**C**OMPUTER **T**OMOGRAPHY:  
xrays absorbed



<http://science.howstuffworks.com/nuclear-medicine1.htm>

**P**OSITRON **E**MISSION **T**OMOGRAPHY:  
positrons emitted

*central serous chorioretinopathy*






<http://vsri.ucdavis.edu/research/retinal/oct>

**O**PTICAL **C**OHERENCE **T**OMOGRAPHY:  
coherent light

# imaging (metabolic): positron emission tomography (PET)

- insert radioactive tracer into blood stream  
e.g. 2-deoxy-2-( $^{18}\text{F}$ )fluoro-D-glucose (FDG)
- FDG (*viz* 'glucose' and metabolism) concentrates in areas of high neural activity = high metabolism
- $^{18}\text{F}$  isotope emits positrons ('electrons' with + charge)
- detectors map areas of positron emission



<http://healthyscientist.blogspot.com/2009/11/wonders-of-pet-scan.html>

<http://www.drugabuse.gov/publications/teaching-packets/understanding-drug-abuse-addiction/section-ii/2-positron-emission-tomography-pet-scan-person-us>

## **THE BOLD SIGNAL : BLOOD OXYGEN LEVEL DEPENDENT fMRI**

- Magnetic Resonance Imaging involves putting the sample (*i.e. your head!!*) in a large external magnet
- Brain imaging is most often done with a spectrometer that measures the magnetic properties of hydrogen (in water) in various parts of the brain
- The hydrogen nucleus behaves like a tiny magnet with different energies when aligned with or against an external magnet. However the exact energies also depend on nearby molecules that provide 'local' magnetic fields. The 'flip' energies are measured using radio wave pulses.
- (via the hemoglobin molecule) Deoxygenated blood is magnetic while oxygenated blood is not magnetic (TAKE CHEM 1B!!).


Also See: [What is Functional Magnetic Resonance Imaging \(fMRI\)?, Hannah Devlin](#)

[fMRI an Introduction: Michael Firbank](#)



## THE **BOLD** SIGNAL : **B**LOOD**O**XYGEN**L**EVEL**D**EPENDENT fMRI (cont.)

- Near inactive neurons there is both oxygenated and deoxygenated blood due to normal metabolic activity
- When a neuron fires there is a momentary increase in the nearby deoxygenated (magnetic) blood but then a rush of blood flow brings fresh a great concentration of oxygenated (nonmagnetic) blood in nearby capillaries
- fMRI monitors the **BOLD** signal for parts of the brain with active vs inactive neurons




<http://psychcentral.com/lib/what-is-functional-magnetic-resonance-imaging-fmri/>

### fMRI

#### *BOLD and Haemodynamic effects*

- Increased energy consumption extracts oxygen
- Vasodilation and CBF increase oxygen supply



- Hemodynamic response time of ~3s

<http://www.slideshare.net/christina101/fmri-introduction-newcastle-university-newcastle-upon-tyne>

# imaging (metabolic): functional Magnetic Resonance Imaging (fMRI)



## The neural correlates of maternal and romantic love





Fig. 3. Overlap between activity of maternal love and romantic love. Activity obtained in this study (contrast: cO vs. cA) was colored in yellow and overlaid on sections through a template brain, along with activity obtained in our previous study on romantic l...

Andreas Bartels, Semir Zeki

NeuroImage, Volume 21, Issue 3, 2004, 1155–1166

## imaging (binding agents): $Ca^{2+}$ imaging



- the flow of  $Ca^{2+}$  ions is associated with neuronal activity (e.g. synaptic vesicles, *remember !!*)
- certain molecules that change their fluorescence (“glow”) when attached to  $Ca^{2+}$
- these molecules are inserted into neurons
- the brain preparation is excited with UV light and the neurons fluorescence (“glow”) with an intensity depending on concentration of  $Ca^{2+}$
- cameras used to follow the action potentials fluorescence as  $Ca^{2+}$  ‘flows’



**Imaging Calcium in Neurons**  
Christine Grienberger  
Arthur Konnerth  
Neuron 73, pp 862-885 (2012)

# functional imaging: voltage sensitive dyes

- certain dye molecules change their optical properties [for example the color that they 'glow' fluoresce)] when in the strong electric field of an action potential
- these dyes are painted on the surface of the cortex
- cortical areas with neurons responding to a specific stimulus fluoresce with different wavelengths (colors) than do inactive areas



<http://www.nature.com/nrn/journal/v5/n11/full/nrn1536.html>

### 3. Imaging

- ✓ a. Positron emission tomography (PET)
- ✓ b. Functional magnetic resonance imaging (fMRI)
- ✓ c. Calcium dyes
- ✓ d. Voltage sensitive dyes

4. Neural activation by external stimulus
  - a. Optogenetics
  - b. Intracranial electrical stimulation
  - c. Transcranial magnetic stimulation (TMS)


- sensory input (light, sound, rat whiskers !!)
- pharmacology (drugs, etc.)
- stimulating electrode (as in neuron lectures)
- **optogenetics**
- **intracranial electrical stimulation**
- **transcranial magnetic stimulation**

- **genetically modify neurons to express LIGHT SENSITIVE ion channels e.g. Na<sup>+</sup> or Cl<sup>-</sup> channels**
- **activate neuron (depolarize or hyperpolarize) by light source (e.g. laser)**
- **image subsequent neural activity (e.g. Ca<sup>2+</sup> imaging) or behavior in awake animal**



# neural excitation by external stimulus: optogenetics

- genetically modify neurons to express **LIGHT SENSITIVE** ion channels  
e.g.  $\text{Na}^+$  or  $\text{Cl}^-$  channels



**Channelrhodopsin-2 (ChR2) and halorhodopsin (NpHR) and light-sensitive proteins used in optogenetic studies.** ChR2 is a cation channel that opens in response to blue light, whereas NpHR is a chloride pump that is activated by yellow light.

<http://sites.sinauer.com/psychopharm2e/webbox09.03.html>




Fig 1a: Schematic representation of the action of Channelrhodopsin2 and Halorhodopsin on neural cells.





Fig 1b: Activation of action potentials by ChR2 (blue light) and inactivation by NpHR (yellow light) of a cultured hippocampal cell

<http://optogenetics.weebly.com/why--how.html>

# neural excitation by external stimulus: optogenetics




---

- genetically modify neurons to express **LIGHT SENSITIVE** ion channels  
e.g. Na<sup>+</sup> or Cl<sup>-</sup> channels



# neural excitation by external stimulus: optogenetics

- activate neuron (depolarize or hyperpolarize) by light source (e.g. laser)




<http://optogenetics.weebly.com/why--how.html>

<http://sciencecalling.com/2011/08/12/light-science-optogenetics/>


<http://web.stanford.edu/group/dlab/optogenetics/>

# neural excitation by external stimulus: optogenetics

- image subsequent neural activity (e.g.  $\text{Ca}^{2+}$  imaging) or behavior in awake animal




<http://nmbi.stanford.edu/research/experiments/optogenetics.htm>




<http://sites.sinauer.com/psychopharm2e/webbox09.03.html>

**Neurons expressing ChR2 or NpHR can be exposed to light of the appropriate wavelength** while the animal is behaving freely in an apparatus such as an operant chamber. Performing an operant response such as a nose-poke turns a laser light generator on for a short period of time to activate light-sensitive proteins in the vicinity of a previously implanted optic fiber

## Method of the Year 2010



<http://www.nature.com/nmeth/video/moy2010/index.html>



<http://www.jove.com/video/51483/in-vivo-optogenetic-stimulation-of-the-rodent-central-nervous-system>

# optogenetics: Prof. Kliger interview



UNIVERSITY OF CALIFORNIA  
SANTA CRUZ

NEWSCENTER

## Ion channel mechanics yield insights into optogenetics experiments

July 06, 2015  
By Tim Stephens

SHARE THIS STORY: [t](#) [f](#) [g+](#) [in](#) [v](#)

Optogenetics techniques, which allow scientists to map and control nerve cells using light stimulation, are being used to study neural circuits in the brain with unprecedented precision. This revolutionary technology relies on light-sensitive proteins such as channelrhodopsins, and researchers at UC Santa Cruz have now determined the molecular mechanism involved in the light-induced activation of one of these proteins.



The new findings, published July 3 in two papers in the *Journal of Biological Chemistry*, can help scientists create tailor-made proteins optimized for use in optogenetics, said David Kliger, senior author of both papers and a professor of chemistry and biochemistry at UC Santa Cruz.

"Little was known about the functional mechanism of these proteins even though they are widely used in optogenetics," Kliger said.


The researchers used fast laser spectroscopy to study the function of Channelrhodopsin-2, which is found in a type of marine algae and is widely used in optogenetics experiments. Channelrhodopsins are ion channels that control the flow of ions across cell membranes. There are many kinds of ion channels that serve different purposes in different types of cells. Nerve signals involve ion flow across the



David Kliger's lab uses fast laser spectroscopy to study biologically important molecules such as channelrhodopsin, an ion channel widely used in optogenetics experiments.



## Crown 85- Visual Perception: A Window to Brain and Behavior



HOME COURSE INFORMATION VISION LECTURES REPORTS RESEARCH LAB INTERVIEWS PROJECTS eCOMMONS

One goal of Crown 85 is to introduce its students to researchers and research in the various neuroscience and perception laboratories throughout the UCSC campus. Enrolled students are required to visit one of these researchers and present a class report on the studies being conducted in their laboratory. Laboratories which have agreed to host Crown 85 student interviews include:

Prof. David Kliger  
Department of Chemistry and  
Biochemistry

Structure and Spectroscopy of  
Visual Pigments  
Early events in Visual  
Processing

[read more](#)




<http://news.ucsc.edu/2015/07/optogenetics.html>

- *neural excitation by external stimulus: intracranial electrical stimulation*

---

**intracranial electrical stimulation:** direct electrical stimulation of brain in awake subjects either with temporary or implanted electrodes (in consenting patients often those with epilepsy) in order to:

- map brain areas to guide surgical procedures
- to monitor brain function in patients
- to explore cognitive responses




<http://golbylab.bwh.harvard.edu/intracranialEEG/EEG.html>

- *neural excitation by external stimulus: intracranial electrical stimulation*

## Electrical Stimulation of the Left and Right Human Fusiform Gyrus Causes Different Effects in Conscious Face Perception

Vinitha Rangarajan<sup>1,2</sup>, Dora Hermes<sup>2</sup>, Brett L. Foster<sup>1,2</sup>, Kevin S. Weiner<sup>2,3</sup>, Corentin Jacques<sup>2,3,4</sup>, Kalanit Grill-Spector<sup>2,3,5</sup>, and Josef Parvizi<sup>1,2,5</sup>




<http://www.jneurosci.org/content/34/38/12828.long>

Bilateral face-selective ECoG responses in the fusiform gyrus. **a**, Face-selective HFB responses in the FG were measured bilaterally




- **transcranial magnetic stimulation (TMS)**

- external magnetic coil put on head, pulses direct magnetic field to stimulate brain areas



First developed in 1985, rTMS has been studied as a possible treatment for **depression**, **psychosis** and other disorders since the mid-1990's. Clinical trials studying the effectiveness of rTMS reveal **mixed results**. When compared to a placebo or inactive (sham) treatment, some studies have found that rTMS is more effective in treating patients with major depression but other studies have found no difference in response compared to inactive treatment.

## repetitive transcranial magnetic stimulation (rTMS) reduces activity



Nature Reviews | Neuroscience



b | Brain images from a study that used positron emission tomography (PET) to measure metabolic activity. The colour coding shows the areas in which activity after a 25 min session of real 1-Hz is **less than that seen after a sham rTMS session. There are significant decreases in activity after real rTMS at the site of stimulation (outlined in red) as well as at many distant sites.** L, left side of the brain.

4. Neural activation by external stimulus
  - ✓ a. Optogenetics
  - ✓ b. Intracranial electrical stimulation
  - ✓ c. Transcranial magnetic stimulation (TMS)

**Psychophysics** is the scientific **study** of the relationship between stimuli (specified in physical terms) and the sensations and perceptions evoked by these stimuli. The term **psychophysics** is used to denote both the substantive **study** of stimulus-response relationships and the methodologies used for this **study**.

[http://www.cis.rit.edu/people/faculty/montag/vandplite/pages/chap\\_1/ch1p2.html](http://www.cis.rit.edu/people/faculty/montag/vandplite/pages/chap_1/ch1p2.html)

# psychophysics in the switkes group



$Threshold = 0.294957 (\pm 0.00499865)$   
 $\sigma = 0.05350 (\pm 0.00470472)$   
 $\chi^2 = 1.53651$   
 95% Fiducial Limits: 0.305464, 0.284777



## *comparison of attributes of some brain recording techniques*

---

	<b>fast or slow</b>	<b>resolution</b>	<b>local or global</b>	<b>Invasive nonInvasive</b>
Single cell recording	fast	high	local	invasive
Electroencephalography (EEG)	fast	low	global	noninvasive
Magnetoencephalography (MEG)	fast	moderate	global	noninvasive
Positron emission tomography (PET)	slow	low	global	noninvasive (but involves radioactive material)
fMRI	slow	low	global	noninvasive
Ca <sup>2+</sup> dyes	fast	high	intermediate	invasive
optogenetics	fast	high	intermediate	invasive

what types of neural processes would each of these be suited to measure ?

*Finis*